

Product datasheet for MC212172

U2af1 (NM_024187) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: U2af1 (NM_024187) Mouse Untagged Clone

Tag: Tag Free Symbol: U2af1

Synonyms: 35kDa; 2010107D16Rik

Mammalian Cell N

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC212172 representing NM_024187

Red=Cloning site Blue=ORF

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul ACCN: NM 024187

Insert Size: 720 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

> point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: BC115479, AAI15480

RefSeq Size: 849 bp RefSeq ORF: 720 bp Locus ID: 108121 **UniProt ID:** Q9D883

17 B1 Cytogenetics:

Gene Summary: Plays a critical role in both constitutive and enhancer-dependent splicing by mediating

> protein-protein interactions and protein-RNA interactions required for accurate 3'-splice site selection. Recruits U2 snRNP to the branch point. Directly mediates interactions between U2AF2 and proteins bound to the enhancers and thus may function as a bridge between U2AF2 and the enhancer complex to recruit it to the adjacent intron (By similarity).

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) encodes isoform 1. Variants 1 and 2 are the same length, as

are isoforms 1 and 2; however, their sequences vary slightly.