OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for MC212451

Pirt (NM_178656) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pirt (NM_178656) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Pirt
Synonyms:	A530088H08Rik
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	SgfI-MluI
ACCN:	NM_178656
Insert Size:	408 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_178656.3, NP_848771.1

View online »

RefSeq Size:	3014 bp
RefSeq ORF:	408 bp
Locus ID:	193003
UniProt ID:	Q8BFY0
Cytogenetics:	11 B3
Gene Summary:	Regulatory subunit of TRPV1, a molecular sensor of noxious heat and capsaicin. Positively regulates TRPV1 channel activity via phosphatidylinositol 4,5-bisphosphate (PIP2). Binds various phosphoinositide, including phosphatidylinositol 4,5-bisphosphate (PIP2), but not phosphatidylinositol (PI).[UniProtKB/Swiss-Prot Function]