

## Product datasheet for MC214381

### H2ac10 (NM\_175661) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids

**Tag:** Tag Free

**Symbol:** H2ac10

**Synonyms:** H2a-22; H2a-22l; Hist1h; Hist1h2af

**Mammalian Cell Selection:** Neomycin

**Vector:** pCMV6-Entry (PS100001)

**E. coli Selection:** Kanamycin (25 ug/mL)

**Fully Sequenced ORF:** >MC214381 representing NM\_175661  
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCCGCGATCGCC

ATGTCTGGACGCGGCAACAGGGTGGCAAGGCTCGCGCAAGGCCAAGACCCGCTCCTCCCGGGCCGGCC  
TGCAGTTCCTCCGCTGGGCGCGTGCACCGGCTGCTCCGCAAGGGCAACTACTCGGAGCGCGTGGGCGCCGG  
CGCCCCGGTGTACCTGGCGGCCGTGCTGGAGTACCTGACGGCCGAGATCCTGGAGCTGGCGGGCAACGCG  
GCCCGCGACAACAAGAAGACGCGCATCATCCGCGCCACCTGCAGCTGGCCATCCGCAACGACGAGGAGC  
TCAACAAGCTGTTGGGCGCGTGACCATCGCGCAGGGCGGCGTCTGCCCAACATCCAGGCCGTGCTGCT  
GCCCAAGAAGACCGAGAGCCACCATAAGCCCAAGGAAAGTAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-MluI

**ACCN:** NM\_175661

**Insert Size:** 393 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



|                               |  |
|-------------------------------|--|
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>  |
| <b>Note:</b>                  | Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.   |
| <b>RefSeq:</b>                | <u>NM_175661.2, NP_783592.1</u>  |
| <b>RefSeq Size:</b>           | 468 bp   |
| <b>RefSeq ORF:</b>            | 393 bp   |
| <b>Locus ID:</b>              | 319173   |
| <b>UniProt ID:</b>            | <u>Q8CGP5</u>  |
| <b>Cytogenetics:</b>          | 13 A3.1  |
| <b>Gene Summary:</b>          | <p>Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. [provided by RefSeq, Aug 2015]</p> |