

## Product datasheet for MC215799

## Sqstm1 (NM\_011018) Mouse Untagged Clone

## **Product data:**

Product Type:	Expression Plasmids
Product Name:	Sqstm1 (NM_011018) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Sqstm1
Synonyms:	A170; OSF-6; Osi; p62; STAP; STONE14
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Sqstm1 (NM_011018) Mouse Untagged Clone – MC215799	
Fully Sequenced ORF:	>NM_011018.2 Red=Cloning site Blue=ORF Green=Tags(s)
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCGTCGTTCACGGTGAAGGCCTATCTTCTGGGCAAGGAGGAGGCGACCCGCGAGATCCGCCGCTTCA GCTTCTGCTTCAGCCCGAGCCGAG
	ACCAAGAATTACGACATCGGGGCTGCTCTGGACACGATCCAGTATTCGAAGCACCCTCCACCATTGTGA ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAGGTTTAA
Chromatograms:	https://cdn.origene.com/chromatograms/ja1382_e02.zip
<b>Restriction Sites:</b>	Sgfl-Mlul
ACCN:	NM_011018
Insert Size:	1329 bp
OTI Disclaimer:	Due to the inherent nature of this plasmid, standard methods to replicate additional ar of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <u>custsupport@origene.com</u> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

<b>ORIGENE</b> Sqstm1 (NM_011018) Mouse Untagged Clone – MC215799	
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 011018.2, NP 035148.1</u>
RefSeq Size:	2000 bp
RefSeq ORF:	1329 bp
Locus ID:	18412
UniProt ID:	<u>Q64337</u>
Cytogenetics:	11 B1.3
Gene Summary:	Autophagy receptor required for selective macroautophagy (aggrephagy). Functions as a bridge between polyubiquitinated cargo and autophagosomes. Interacts directly with both

the cargo to become degraded and an autophagy modifier of the MAP1 LC3 family. Required both for the formation and autophagic degradation of polyubiquitin-containing bodies, called ALIS (aggresome-like induced structures) and links ALIS to the autophagic machinery. Involved in midbody ring degradation (By similarity). May regulate the activation of NFKB1 by TNF-alpha, nerve growth factor (NGF) and interleukin-1. May play a role in titin/TTN downstream signaling in muscle cells. May regulate signaling cascades through ubiquitination. Adapter that mediates the interaction between TRAF6 and CYLD (PubMed:14960283, PubMed:18382763). May be involved in cell differentiation, apoptosis, immune response and regulation of K(+) channels. Involved in endosome organization by retaining vesicles in the perinuclear cloud: following ubiquitination by RNF26, attracts specific vesicle-associated adapters, forming a molecular bridge that restrains cognate vesicles in the perinuclear region and organizes the endosomal pathway for efficient cargo transport (By similarity). Promotes relocalization of 'Lys-63'-linked ubiquitinated TMEM173/STING to autophagosomes (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (1) represents the longer transcript and encodes the longer isoform (1).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US