

## **Product datasheet for MC215996**

## Cir1 (NM\_025854) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Cir1 (NM\_025854) Mouse Untagged Clone

Tag: Tag Free

Symbol: Cir1

**Synonyms:** 1700023B02Rik; 2810021A19Rik; Cicr; Cir

Mammalian Cell

Selection:

Neomycin

**Vector:** pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_025854

**Insert Size:** 1353 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 025854.3</u>, <u>NP 080130.2</u>

RefSeq Size: 1945 bp RefSeq ORF: 1353 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Locus ID:** 66935

ORIGENE

UniProt ID: Q9DA19
Cytogenetics: 2 C3

**Gene Summary:** Regulates transcription and acts as corepressor for RBPJ. Recruits RBPJ to the Sin3-histone

deacetylase complex (HDAC). Required for RBPJ-mediated repression of transcription (By similarity). May modulate splice site selection during alternative splicing of pre-mRNAs.

[UniProtKB/Swiss-Prot Function]