

Product datasheet for **MC216088**

Pold3 (NM_133692) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pold3 (NM_133692) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Pold3
Synonyms:	2410142G14Rik; C85233; P66; P68
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC216088 representing NM_133692
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGC**C

ATGGCGGAACAGCTGTATCTAGAAAACATAGACGAGTTCGTACGGACCAGAACAAGATCGTGACTTACA
 AGTGGCTAAGCTATACACTAGGGGTTTCATGTTAACAGGCCAAAACAGATGCTCTATGAATATGTTGAAAG
 GAAACGGAAAGAAAATTCGGGAGCTCAGCTGCATGTTACCTACTTGGTGTCTGGCAGTCTTATACAGAAC
 GGACATTTCTGCCACAAGTTGCAGTAGTGAGAGAAGATAAACTGGAAGCAGTGAAGTCCAAGTTAGCTG
 TGACTGCCAGCATCCATGTGTACAGCATCCAGAAAGCTATGCTAAAGGACAGTGGGCCTCTGTTCAATAC
 CGACTATGACATCCTTAAAGCAATTTGCAGAACTGCAGCAAGTTTGTGCCATACAGTGTGCAGCTGCA
 GTCCCCAGAGCTCCTGCAGAAATCCCATCTTCCAGAAAGTATGAACAGTCAAATCTTCAGGCAGCGAGTG
 AGGCACAAGCCAGTGAGCTGACTACCAATGGCCATGGTCCACCTGCCTCCAAACAGGCTTCCCAGCAGCC
 CAAAGGAATTATGGGAATGTTAATCTCTAAAGCTGCTACTAAAACCCAGGACACCAACAAGGAAACAAA
 CCAGAGGGCCCGAGAAGTAACAGTGCATCTTCTGCTGGGGCAAAGCACCAGGAAAAGGGAGTGTGATGA
 GCAACTTTTTTGGAAAAGCTGCAATGAATAAACTTAAAGTCAATTTGGATTTCAGAACAAAGCAGTAAAGGA
 AGAAAAACAGTGGAGCAACCTCCAGTGTCTGTCACTGAACCAAAGCTGGCAGCTCCCCAGCTCAGAAG
 AAATCCAGCAGAAAGTCCGAGCCTGGGAAGGTGCAGCAGAAAGGAGAAAAGCAGGGGCAAGCGAGTAGACT
 TGTCCGATGAGGAGGCAAAGGAAACCGAACCTGAAGAAAAAGAGAAGAAGAATCAAGTTCCTCAGTC
 TGATAGCAGTGAAGATGAAGTCTTGAAGACTCCCCTGAGATGTATGAAGCAGACTCACCATCTCCACCT
 CCTGTATCTCCACCTCCTGATCCTATGCCAAAACTGAGCCCCCTCCTGTCAAGCGTTCAAGTGGAGAAA
 CCAAAGGAGACGAAAGCGTGTACTGAAATCTAAAACCTTTGTGGATGAAGAAGGCTGCATAGTACTGA
 GAAAGTCTATGAGAGTGAGTCTCTGCACAGACAGTGAGGAGGAGCTTAAGATGAAGCCGGCCTCAGCACAC
 AAACCCCTGCTGCCGCTGTGAAAAGGGAGCCAGAGAAGAACGGAAGGGCCCAAGAAAGGGGCTGCTG
 CTCTGGGCAAAGCCAACAGACAAGTGTCCACTACTGGCTTCTTCCAGAAAAAG**TAA**

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_133692
- Insert Size:** 1386 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_133692.2](#), [NP_598453.1](#)

RefSeq Size: 3002 bp

RefSeq ORF: 1386 bp

Locus ID: 67967

Cytogenetics: 7 E2

Gene Summary: As a component of the trimeric and tetrameric DNA polymerase delta complexes (Pol-delta3 and Pol-delta4, respectively), plays a role in high fidelity genome replication, including in lagging strand synthesis, and repair (PubMed:10219083, PubMed:27524497). Required for optimal Pol-delta activity. Stabilizes the Pol-delta complex and plays a major role in Pol-delta stimulation by PCNA. Pol-delta3 and Pol-delta4 are characterized by the absence or the presence of POLD4. They exhibit differences in catalytic activity. Most notably, Pol-delta3 shows higher proofreading activity than Pol-delta4. Although both Pol-delta3 and Pol-delta4 process Okazaki fragments in vitro, Pol-delta3 may also be better suited to fulfill this task, exhibiting near-absence of strand displacement activity compared to Pol-delta4 and stalling on encounter with the 5'-blocking oligonucleotides. Pol-delta3 idling process may avoid the formation of a gap, while maintaining a nick that can be readily ligated. Along with DNA polymerase kappa, DNA polymerase delta carries out approximately half of nucleotide excision repair (NER) synthesis following UV irradiation. In this context, POLD3, along with PCNA and RFC1-replication factor C complex, is required to recruit POLD1, the catalytic subunit of the polymerase delta complex, to DNA damage sites. Under conditions of DNA replication stress, required for the repair of broken replication forks through break-induced replication (BIR). Involved in the translesion synthesis (TLS) of templates carrying O6-methylguanine or abasic sites performed by Pol-delta4, independently of DNA polymerase zeta (REV3L) or eta (POLH). Facilitates abasic site bypass by DNA polymerase delta by promoting extension from the nucleotide inserted opposite the lesion. Also involved in TLS, as a component of the POLZ complex. Along with POLD2, dramatically increases the efficiency and processivity of DNA synthesis of the minimal DNA polymerase zeta complex, consisting of only REV3L and REV7 (By similarity).[UniProtKB/Swiss-Prot Function]