

## Product datasheet for **MC216139**

### Esrrb (NM\_011934) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Esrrb (NM_011934) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Esrrb
Synonyms:	Err2; Errb; Estrrb; Nr3b2
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

**Fully Sequenced ORF:** >MC216139 representing NM\_011934  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGGACGTGTCCGAACCTGTCATCCCGACCCCTTGGCTACCACAACCAGCTGTGAACCGAATGTCGT  
 CCGAAGACAGGCACCTGGGCTCTAGTTGCGGCTCCTTCATCAAGACGAGCCATCCAGCCGTCCTCGGG  
 CATTGATGCCCTCAGCCACCACAGCCCCAGCGGCTCGTCGGACGCCAGTGGTGGCTTTGGCATTGCCCTG  
 AGCACCCACGCCAACGGTCTGGACTCGCCGCTATGTTTCGACAGTGGGGGCTGGGAGGCAACCCGTGCC  
 GCAAGAGCTACGAGGACTGTACTAGTGGTATCATGGAGGACTCCGCCATCAAATGCGAGTACATGCTTAA  
 CGCCATCCCCAAGCGCTGTGCCTCGTGTGCGGGGACATTGCCTCTGGCTACCACTACGGAGTGGCTCC  
 TGCGAGGCTTGAAGGCTTCTCAAGAGAACCATTCAAGGCAACATCGAGTACAACCTGCCCGGCCACCA  
 ATGAATGTGAGATCACCAAACGGAGGCGCAAGTCTGTGACGGCTGCCGATTCATGAAATGCCTCAAAGT  
 GGGGATGCTGAAGGAAGGTGTGCGCCTTGACCGAGTTCGAGGAGCGCCAGAAGTACAAGCGACGGCTG  
 GATTCGGAGAACAGCCCTACCTGAACCTGCCGATTTCCCAACCTGCTAAAAAGCCATTGACTAAGATCG  
 TCTCGAATCTACTAGGGTTGAGCAGGACAAGCTGTATGCTATGCCTCCCAACGATATCCCCGAGGGAGA  
 TATCAAGGCCCTGACCACTCTCTGTGAATTGGCAGATCGGGAGCTTGTGTTCTCATCAACTGGGCCAAG  
 CACATCCCAGGCTTCCCCAGTCTGACACTTGGGGACCAGATGAGCCTGCTGCAGAGTGCCTGGATGGAGA  
 TTCTCATCTTGGGCATCGTGTACCGCTCGCTCCATACGATGACAAGCTGGCATAACGCCGAGGACTATAT  
 CATGGATGAGGAACACTCTCGCCTGGTAGGGTGTGACCTTTACCGAGCCATCCTGCAGCTGGTGCCG  
 AGGTACAAGAACTCAAGGTAGAGAAGGAAGAGTTTATGATCCTCAAGGCCCTGGCCCTCGCCAACCTCAG  
 ATTCGATGTACATTGAGAACCTGGAGCGGTGCAGAAGTCCAGGACCTGCTGCACGAGGCGCTGCAGGA  
 CTATGAGCTGAGTCAGCGCCACGAGGAGCGCGGAGGGCCGCAAGCTGCTGCTGACGCTGCCCTGCTG  
 AGGCAGACAGCCGCAAGCCGTGCAACACTTCTACAGTGTGAACTGCAGGGCAAGGTGCCCATGCACA  
 AACTCTTCTGGAGATGCTGGAGCCAAGGTG**TGA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM\_011934
- Insert Size:** 1365 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM\\_011934.4](#), [NP\\_036064.3](#)

RefSeq Size: 4208 bp

RefSeq ORF: 1365 bp

Locus ID: 26380

UniProt ID: [Q61539](#)

Cytogenetics: 12 40.49 cM

**Gene Summary:** Transcription factor that binds a canonical ESRRB recognition (ERRE) sequence 5'TCAAGGTCA-3' localized on promoter and enhancer of targets genes regulating their expression or their transcriptional activity (PubMed:27601327, PubMed:23169531, PubMed:23508100, PubMed:26206133, PubMed:20534447, PubMed:18662995, PubMed:18957414, PubMed:27723719, PubMed:23019124). Plays a role, in a LIF-independent manner, in maintenance of self-renewal and pluripotency of embryonic and trophoblast stem cells through different signaling pathways including FGF signaling pathway and Wnt signaling pathways (PubMed:18957414, PubMed:26206133, PubMed:20534447, PubMed:23040478, PubMed:23040477, PubMed:23019124, PubMed:23169531). Upon FGF signaling pathway activation, interacts with KDM1A by directly binding to enhancer site of ELF5 and EOMES and activating their transcription leading to self-renewal of trophoblast stem cells (PubMed:26206133). Also regulates expression of multiple rod-specific genes and is required for survival of this cell type (PubMed:20534447). Plays a role as transcription factor activator of GATA6, NR0B1, POU5F1 and PERM1 (PubMed:18662995, PubMed:23508100, PubMed:18957414). Plays a role as transcription factor repressor of NFE2L2 transcriptional activity and ESR1 transcriptional activity (By similarity). During mitosis remains bound to a subset of interphase target genes, including pluripotency regulators, through the canonical ESRRB recognition (ERRE) sequence, leading to their transcriptional activation in early G1 phase (PubMed:27723719). Can coassemble on structured DNA elements with other transcription factors like SOX2, POU5F1, KDM1A and NCOA3 to trigger ESRRB-dependent gene activation (PubMed:23019124, PubMed:23169531, PubMed:18662995, PubMed:26206133). This mechanism, in the case of SOX2 corecruitment prevents the embryonic stem cells (ESCs) to epiblast stem cells (EpiSC) transition through positive regulation of NR0B1 that inhibits the EpiSC transcriptional program (PubMed:23169531). Also plays a role inner ear development by controlling expression of ion channels and transporters and in early placentation (PubMed:9285590, PubMed:17765677).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) represents the shorter transcript but encodes the longer isoform (1). CCDS Note: An upstream translation start codon is selected for this CCDS for consistency with other family members. The use of an alternative upstream start codon would result in a protein that is 21 aa longer at the N-terminal.