

Product datasheet for **MC216258**

Foxa1 (NM_008259) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Foxa1 (NM_008259) Mouse Untagged Clone

Tag: Tag Free

Symbol: Foxa1

Synonyms: Hnf-3a; Hnf3a; Tcf-3a; Tcf3a

Mammalian Cell Selection: Neomycin

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul

ACCN: NM_008259

Insert Size: 1407 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_008259.3](#), [NP_032285.2](#)

RefSeq Size: 3188 bp

RefSeq ORF: 1407 bp



[View online »](#)

Locus ID: 15375

UniProt ID: [P35582](#)

Cytogenetics: 12 24.7 cM

Gene Summary: Transcription factor that is involved in embryonic development, establishment of tissue-specific gene expression and regulation of gene expression in differentiated tissues. Is thought to act as a 'pioneer' factor opening the compacted chromatin for other proteins through interactions with nucleosomal core histones and thereby replacing linker histones at target enhancer and/or promoter sites. Binds DNA with the consensus sequence 5'-[AC]A[AT]T[AG]TT[GT][AG][CT]T[CT]-3' (By similarity). Proposed to play a role in translating the epigenetic signatures into cell type-specific enhancer-driven transcriptional programs. Involved in the development of multiple endoderm-derived organ systems such as the liver, pancreas, lungs and prostate; FOXA1 and FOXA2 seem to have at least in part redundant roles. Plays a role in prostate morphogenesis and epithelial cell differentiation. FOXA1 and FOXA2 are essential for hepatic specification. FOXA1 and FOXA2 are required for morphogenesis and cell differentiation during formation of the lung. FOXA1 and FOXA2 are involved in bile duct formation; they positively regulate the binding of glucocorticoid receptor/NR3C1 to the IL6 promoter. FOXA1 and FOXA2 regulate multiple phases of midbrain dopaminergic neuron development; they regulate expression of NEUROG2 at the beginning of mDA neurogenesis and of NR4A2 and EN1 in immature mDA neurons. Modulates the transcriptional activity of nuclear hormone receptors. Is involved in ESR1-mediated transcription. Inhibits NKX2-1-mediated transcription from the SFTPC promoter in lung epithel independently from DNA-binding. Involved in regulation of apoptosis. Involved in cell cycle regulation. Originally described as a transcription activator for a number of liver genes such as AFP, albumin, tyrosine aminotransferase, PEPCK, etc. Interacts with the cis-acting regulatory regions of these genes. Involved in glucose homeostasis; activates the GCG promoter.[UniProtKB/Swiss-Prot Function]