

Product datasheet for **MC216418**

Nono (NM_023144) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Nono (NM_023144) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Nono
Synonyms:	AA407051; AV149256; nonA; NRB54; P54NRB
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC216418 representing NM_023144
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGGATCGCC**

ATGCAGAGCAATAAAGCCTTTAACTTGGAGAAGCAGAATCATACTCCAAGGAAGCATCATCAGCATCACC
ACCAGCAGCACCATCAGCAGCAACAGCAGCAGCAGCAACAGCCACCCCAACATACCTGCAAATGG
CCAGCAGGCCAGCAGCCAGAATGAAGGCTTGACTATTGACCTGAAGAATTTAGGAAACCAGGAGAGAAG
ACCTTTACACAGCGTAGCCGTCTCTTTGTGGGCAATCTTCCCCTGATATCACTGAGGAGGAAATGAGGA
AACTATTTGAGAAATATGGAAAAGCAGGCGAAGTTTTCATTTCATAAGGATAAAGGCTTTGGCTTTATTTCG
CTTGAAAACACGAACCTAGCGGAAATGCCAAAGTGGAGCTGGACAACATGCCCTCCGTGGGAAGCAG
CTGCGAGTGCCTTTGCCTGTACAGTGCATCCCTTACAGTCCGCAACCTTCCTCAGTACGTGTCGAACG
AACTGCTGGAAGAAGCCTTTTCTGTGTTGCGCCAGGTGGAGGGCTGTAGTCATTGTGGATGACCGAGG
AAGGCCCTCAGGAAAGGCATTGTTGAGTTCTCAGGGAAGCCAGCTGCTCGGAAAGCTCTGGACAGATGC
AGTGAAGGCTCCTTCTGCTGACTACATTTCTCGGCCTGTGACTGTGGAGCCTATGGACCAGTTAGATG
ATGAAGAGGGACTTCCAGAGAACTGGTTATAAAAAACCAGCAATTCACAAGGAGAGAGAACAGCCACC
CAGATTTGCACAACCTGGCTCCTTTGAGTATGAGTATGCCATGCGCTGGAAGGCACTCATTGAGATGGAG
AAGCAACAGCAGGATCAAGTGGATCGGAACATCAAGGAGGCTCGTGAGAAGCTGGAGATGGAGATGGAGG
CTGCACGTCAAGCACCAGGTTATGCTAATGAGGCAGGATTTGATGAGACGTCAAGAAGAGCTTCGGAG
AATGGAGGAGCTGCATAACCAAGAGGTTTCAAGAGCAAGAGCAAGTGAAGTCAAGCAGGAAAGAGGAAACGC
AGGCGCCGTGAGGAAGAGATGCGGCGACAGCAAGAGGAAATGATGCGCCGACAGCAGGAAGGATTCAGG
GAACCTTCCCTGATGCGAGAGAACAAGAGATACGGATGGGCCAAATGGCTATGGGAGGTGCTATGGGCAT
AAACAATAGAGGCGCGATGCCCTGCTCCTGTGCCACCTGGTACTCCAGCTCCTCCAGGACCTGCCACT
ATGATGCCAGATGGAACCTTGGATTGACCCACCAACAAGTGAACGTTTTGGCCAAAGCTGCAACATGG
AAGGAATTGGAGCAATTGGTGGAACTCCTCCTGCATTCAACCGTCCAGCTCCGGGAGCTGAATTTGCTCC
AAATAACGCCGCCGAT**TAG**

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI

ACCN: NM_023144

Insert Size: 1422 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_023144.2](#), [NP_075633.2](#)

RefSeq Size: 2468 bp

RefSeq ORF: 1422 bp

Locus ID: 53610

UniProt ID: [Q99K48](#)

Cytogenetics: X D

Gene Summary: DNA- and RNA binding protein, involved in several nuclear processes. Binds the conventional octamer sequence in double-stranded DNA. Also binds single-stranded DNA and RNA at a site independent of the duplex site. Involved in pre-mRNA splicing, probably as a heterodimer with SFPQ. Interacts with U5 snRNA, probably by binding to a purine-rich sequence located on the 3' side of U5 snRNA stem 1b. Together with PSPC1, required for the formation of nuclear paraspeckles. The SFPQ-NONO heteromer associated with MATR3 may play a role in nuclear retention of defective RNAs. The SFPQ-NONO heteromer may be involved in DNA unwinding by modulating the function of topoisomerase I/TOP1. The SFPQ-NONO heteromer may be involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination and may stabilize paired DNA ends. In vitro, the complex strongly stimulates DNA end joining, binds directly to the DNA substrates and cooperates with the Ku70/G22P1-Ku80/XRCC5 (Ku) dimer to establish a functional preligation complex. NONO is involved in transcriptional regulation. The SFPQ-NONO-NR5A1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity. NONO binds to an enhancer element in long terminal repeats of endogenous intracisternal A particles (IAPs) and activates transcription. Regulates the circadian clock by repressing the transcriptional activator activity of the CLOCK-ARNTL/BMAL1 heterodimer. Important for the functional organization of GABAergic synapses. Plays a specific and important role in the regulation of synaptic RNAs and GPHN/gephyrin scaffold structure, through the regulation of GABRA2 transcript (PubMed:26571461). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) represents the longer transcript. Variants 1 and 2 encode the same protein.