

Product datasheet for MC216473

Ntrk2 (NM_008745) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Ntrk2 (NM_008745) Mouse Untagged Clone

Tag: Tag Free Symbol: Ntrk2

Synonyms: GP145-TrkB/GP95-TrkB; Tkrb; trk-B; trkB

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn





Fully Sequenced ORF: >MC216473 represe

>MC216473 representing NM_008745 Red=Cloning site Blue=ORF

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCGCCGCGATCGCC

ATGTCGCCCTGGCTGAAGTGGCATGGACCCGCCATGGCGCGCTCTGGGGCTTATGCCTGCTGGTCTTGG GCTTCTGGAGGGCCTCTCTCGCCTGCCCGACGTCCTGCAAATGCAGTTCCGCTAGGATTTGGTGTACTGA GCCTTCTCCAGGCATCGTGGCATTCCCGAGGTTGGAACCTAACAGCGTTGACCCGGAGAACATCACGGAA ATTCTCATTGCAAACCAGAAAAGGCTAGAAATCATCAATGAAGATGACGTTGAAGCTTACGTGGGGCTGA GAAACCTTACAATTGTGGATTCCGGCTTAAAGTTTGTGGCTTACAAAGCGTTTCTGAAAAAACAGCAACCT GCGGCACATAAATTTCACACGAAACAAGCTGACGAGTTTGTCCAGGAGACATTTCCGCCACCTTGACTTG TCTGACCTGATCCTGACGGGTAATCCGTTCACGTGCTCCTGCGACATCATGTGGCTCAAGACTCTCCAGG AGACTAAATCCAGCCCCGACACTCAGGATTTGTACTGCCTCAATGAGAGCAGCAAGAACATGCCCCTGGC GAACCTGCAGATACCCAATTGTGGTCTGCCATCTGCACGTCTGGCTGCTCCTAACCTCACCGTGGAGGAA GGAAAGTCTGTGACCCTTTCCTGCAGTGTGGGGGGTGACCCACTCCCCACCTTGTACTGGGACGTTGGGA TGATGACAGTGGAAAGCAAATCTCTTGTGTGGCAGAAAACCTTGTAGGAGAAGATCAAGATTCTGTGAAC CTCACTGTGCATTTTGCGCCAACTATCACGTTTCTCGAGTCTCCAACCTCAGATCACCACTGGTGCATTC CATTCACTGTGAGAGGCAACCCCAAGCCTGCGCTTCAGTGGTTCTACAATGGGGCCATACTGAATGAGTC CAAGTACATCTGTACTAAGATCCACGTCACCAATCACACGGAGTACCATGGCTGCCTCCAGCTGGATAAC CCCACTCATATGAATAACGGAGACTACACCCTGATGGCCAAGAACGAGTATGGGAAGGATGAGAGACAGA TCTCCGCTCACTTCATGGGCCGGCCTGGAGTCGACTACGAGACAAACCCAAATTACCCTGAAGTCCTCTA TGAAGACTGGACCACGCCAACTGACATTGGGGATACTACGAACAAAGTAATGAAATCCCCTCCACGGAT GTTGCTGACCAAAGCAATCGGGAGCATCTCTCGGTCTATGCCGTGGTGGTGATTGCATCTGTGGTGGGAT TCTGCCTGCTGGTGATGTTGCTCCTGCTCAAGTTGGCGAGACATTCCAAGTTTGGCATGAAAGGTTTTGT TTTGTTTCATAAGATCCCACTGGATGGGTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul
ACCN: NM_008745
Insert Size: 1431 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.



Cytogenetics:

Ntrk2 (NM_008745) Mouse Untagged Clone - MC216473

Plasmids are not sterile. For experiments where strict sterility is required, filtration with Note:

0.22um filter is required.

RefSeq: BC052014, AAH52014

RefSeq Size: 2279 bp RefSeq ORF: 1431 bp Locus ID: 18212 **UniProt ID:** P15209 13 31.2 cM

Gene Summary: Receptor tyrosine kinase involved in the development and the maturation of the central and

> the peripheral nervous systems through regulation of neuron survival, proliferation, migration, differentiation, and synapse formation and plasticity. Receptor for BDNF/brainderived neurotrophic factor and NTF4/neurotrophin-4. Alternatively can also bind

> NTF3/neurotrophin-3 which is less efficient in activating the receptor but regulates neuron

survival through NTRK2. Upon ligand-binding, undergoes homodimerization,

autophosphorylation and activation. Recruits, phosphorylates and/or activates several downstream effectors including SHC1, FRS2, SH2B1, SH2B2 and PLCG1 that regulate distinct overlapping signaling cascades. Through SHC1, FRS2, SH2B1, SH2B2 activates the GRB2-Ras-

MAPK cascade that regulates for instance neuronal differentiation including neurite outgrowth. Through the same effectors controls the Ras-PI3 kinase-AKT1 signaling cascade that mainly regulates growth and survival. Through PLCG1 and the downstream protein kinase C-regulated pathways controls synaptic plasticity. Thereby, plays a role in learning and memory by regulating both short term synaptic function and long-term potentiation. PLCG1

also leads to NF-Kappa-B activation and the transcription of genes involved in cell survival. Hence, it is able to suppress anoikis, the apoptosis resulting from loss of cell-matrix interactions. Isoform GP95-TRKB may also play a role in neutrophin-dependent calcium signaling in glial cells and mediate communication between neurons and glia.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) lacks several exons and uses an alternate 3'-terminal exon, compared to variant 1. The encoded isoform (b) has a shorter and distinct C-terminus, compared to isoform a.