

## **Product datasheet for MC216640**

## Etv4 (NM\_008815) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Etv4 (NM\_008815) Mouse Untagged Clone

Tag: Tag Free

Symbol: Etv4

Synonyms: AW414408; Pea-3; Pea3

Mammalian Cell

Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul

ACCN: NM\_008815

**Insert Size:** 1458 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 008815.3</u>, <u>NP 032841.2</u>

RefSeq Size: 2349 bp RefSeq ORF: 1458 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Etv4 (NM\_008815) Mouse Untagged Clone - MC216640

**Locus ID:** 18612

UniProt ID: P28322

Cytogenetics: 11 65.48 cM

**Gene Summary:** Transcriptional activator that binds to the enhancer of the adenovirus E1A gene; the core-

binding sequence is 5'[AC]GGA[AT]GT-3'. May play a regulatory role during embryogenesis.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) uses an alternate in-frame splice site in the 5' coding

region, compared to variant 1. The encoded isoform (2) is shorter than isoform 1.