

Product datasheet for **MC216904**

Fkrp (NM_173430) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Fkrp (NM_173430) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Fkrp
Synonyms:	A830029B19Rik; AI842067; AI847300; LGMD1I; MDC1C
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_173430
Insert Size:	1485 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_173430.2</u> , <u>NP_775606.1</u>
RefSeq Size:	2817 bp
RefSeq ORF:	1485 bp



[View online »](#)

Locus ID: 243853

UniProt ID: [Q8CG64](#)

Cytogenetics: 7 A2

Gene Summary: Catalyzes the transfer of CDP-ribitol to ribitol 5-phosphate previously attached by FKTN/fukutin of to the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan (DAG1) (By similarity). This constitutes the second step in the formation of the ribose 5-phosphate tandem repeat which links the phosphorylated O-mannosyl trisaccharide to the ligand binding moiety composed of repeats of 3-xylosyl-alpha-1,3-glucuronic acid-beta-1 (By similarity).[UniProtKB/Swiss-Prot Function]