

Product datasheet for **MC217450**

Ceacam1 (NM_001039185) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ceacam1 (NM_001039185) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ceacam1
Synonyms:	bb-1; Bgp; Bgp1; C-CAM; Cc1; CD66a; Cea-1; Cea-7; Cea1; Cea7; Hv-2; Hv2; mCEA1; Mhv-1; MHVR; MHVR1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >MC217450 representing NM_001039185
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGAGCTGGCCTCAGCACATCTCCACAAAGGGCAGGTTCCCTGGGGAGGACTACTGCTCACAGCCTCAC
 TTTTAGCCTCCTGGAGCCCTGCCACCACTGCTGAAGTCACCAATTGAGGCTGTGCCGCCCAGGTTGCTGA
 AGACAACAATGTTCTTCTACTTGTTCACAACTGCCCCCTGGCGCTTGGAGCCTTTGCCTGGTACAAGGGA
 AACACTACGGCTATAGACAAAGAAATTGCACGATTTGTACCAATAGTAATATGAATTTACGGGGCAAG
 CATACAGCGGCAGAGAGATAATATACAGCAATGGATCCCTGCTCTTCAAATGATCACCATGAAGGATAT
 GGGAGTCTACACACTAGATATGACAGATGAAACTATCGTCGACTCAGGCGACTGTGCGATTTTCATGTA
 CACCCCATATTATTAAGCCCAACATCACAAGCAACAACCTCAATCCCGTGGAGGGTGACGACTCCGTAT
 CATTAACCTGTGACTCTTACACTGACCCTGATAATATAAACTACCTGTGGAGCAGAAATGGTGAAAGCCT
 TTCAGAAGGTGACAGGCTGAAGCTGTCTGAGGGCAACAGGACTCTCACTTTACTCAATGTCACGAGGAAT
 GACACAGGACCCTATGTGTGTGAAACCCGGAATCCAGTGAGTGCAACCGAAGTGACCCATTGACGCTGA
 ACATTATCTATGGTCCGGACACCCCGATTATATCCCCCTCAGATATTTATTTGCATCCAGGGTCAAACCT
 CAACCTCTCCTGCCATGCAGCCTCTAACCCACCTGCACAGTACTTTTGGCTTATCAATGAGAAGCCCAT
 GCATCCTCCCAAGAGCTCTTTATCCCCAACATCACTACTAATAATAGCGGAACCTATACCTGCTTCGTCA
 ATAACCTGTGCTACTGGCCTCAGTAGGACCACAGTCAAGAACATTACAGTCCTTGAGCCAGTGACTCAGCC
 CTTCTCCAAGTCACCAACACCACAGTCAAAGAACTAGACTCTGTGACCTGACCTGCTTGTGCAATGAC
 ATTGGAGCCAACATCCAGTGGCTCTTCAATAGCCAGAGTCTTCACTCACAGAGAGAATGACACTCTCCC
 AGAACACAGCATCCTCAGAATAGACCTATTAAAGAGGGAAGATGCCGGCGAGTATCAGTGTGAAATCTC
 GAATCCAGTCAGCGTCAGGAGGAGCAACTCAATCAAGCTGGACATAATATTTGACCCAACACAAGGAGGC
 CTCTCAGATGGCGCCATTGCTGGCATCGTGATTGGAGTTGTGGCTGGGGTGGCTCTAATAGCAGGGCTGG
 CATATTTCTCTATTCCAGGAAGTCTGGCGGGGAAGTGACCAGCGAGATCTCACAGAGCACAAACCTC
 AGCCTCCAACCACAATCTGGCTCCTTCTGACAACCTCTCTAACAAGGTGGATGACGTGCGATACACTGTC
 CTGAACCTCAATCCCAGCAACCAACCGGCCAACTTCAGCCCTTCTTCTCCAAGAGCCACAGAAACAG
 TTTATTGAGAAGTAAAAAGAAG**TGA**

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001039185
Insert Size: 1566 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001039185.1, NP_001034274.1</u>
RefSeq Size:	3736 bp
RefSeq ORF:	1566 bp
Locus ID:	26365
UniProt ID:	<u>P31809</u>
Cytogenetics:	7 13.84 cM
Gene Summary:	<p>Isoform 1: Cell adhesion protein that mediates homophilic cell adhesion in a calcium-independent manner (By similarity). Plays a role as coinhibitory receptor in immune response, insulin action and functions also as an activator during angiogenesis (PubMed:16680193, PubMed:17081782, PubMed:18544705, PubMed:21029969, PubMed:21081647, PubMed:22496641, PubMed:22962327, PubMed:23696226). Its coinhibitory receptor function is phosphorylation- and PTPN6 -dependent, which in turn, suppress signal transduction of associated receptors by dephosphorylation of their downstream effectors (PubMed:17081782, PubMed:21029969, PubMed:22496641). Plays a role in immune response, of T-cells, natural killer (NK) and neutrophils (PubMed:17081782, PubMed:23696226, PubMed:22496641, PubMed:21029969). Upon TCR/CD3 complex stimulation, inhibits TCR-mediated cytotoxicity by blocking granule exocytosis by mediating homophilic binding to adjacent cells, allowing interaction with and phosphorylation by LCK and interaction with the TCR/CD3 complex which recruits PTPN6 resulting in dephosphorylation of CD247 and ZAP70 (PubMed:22496641). Also inhibits T-cell proliferation and cytokine production through inhibition of JNK cascade and plays a crucial role in regulating autoimmunity and anti-tumor immunity by inhibiting T-cell through its interaction with HAVCR2 (PubMed:17081782). Upon natural killer (NK) cells activation, inhibit KLRK1-mediated cytotoxicity of CEACAM1-bearing tumor cells by trans-homophilic interactions with CEACAM1 on the target cell and lead to cis-interaction between CEACAM1 and KLRK1, allowing PTPN6 recruitment and then VAV1 dephosphorylation (PubMed:23696226). Upon neutrophils activation negatively regulates IL1B production by recruiting PTPN6 to a SYK-TLR4-CEACAM1 complex, that dephosphorylates SYK, reducing the production of reactive oxygen species (ROS) and lysosome disruption, which in turn, reduces the activity of the inflammasome (PubMed:22496641). Downregulates neutrophil production by acting as a coinhibitory receptor for CSF3R by downregulating the CSF3R-STAT3 pathway through recruitment of PTPN6 that dephosphorylates CSF3R (PubMed:21029969). Also regulates insulin action by</p>

promoting INS clearance and regulating lipogenesis in liver through regulating insulin signaling (PubMed:18544705). Upon INS stimulation, undergoes phosphorylation by INSR leading to INS clearance by increasing receptor-mediated insulin endocytosis. This internalization promotes interaction with FASN leading to receptor-mediated insulin degradation and to reduction of FASN activity leading to negative regulation of fatty acid synthesis. INSR-mediated phosphorylation also provokes a down-regulation of cell proliferation through SHC1 interaction resulting in decrease coupling of SHC1 to the MAPK3/ERK1-MAPK1/ERK2 and phosphatidylinositol 3-kinase pathways (By similarity). Functions as activator in angiogenesis by promoting blood vessel remodeling through endothelial cell differentiation and migration and in arteriogenesis by increasing the number of collateral arteries and collateral vessel calibers after ischemia (PubMed:16680193, PubMed:22962327). Also regulates vascular permeability through the VEGFR2 signaling pathway resulting in control of nitric oxide production (PubMed:21081647). Downregulates cell growth in response to EGF through its interaction with SHC1 that mediates interaction with EGFR resulting in decrease coupling of SHC1 to the MAPK3/ERK1-MAPK1/ERK2 pathway (PubMed:15467833). Negatively regulates platelet aggregation by decreasing platelet adhesion on type I collagen through the GPVI-FcRgamma complex (PubMed:19008452). Inhibits cell migration and cell scattering through interaction with FLNA; interferes with the interaction of FLNA with RALA (By similarity). Mediates bile acid transport activity in a phosphorylation dependent manner (By similarity). Negatively regulates osteoclastogenesis (PubMed:25490771).[UniProtKB/Swiss-Prot Function]