

## **Product datasheet for MC217756**

## Tfpi (NM\_001177320) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Tfpi (NM\_001177320) Mouse Untagged Clone

Tag: Tag Free
Symbol: Tfpi

Synonyms: A630013F22Rik; AW552122; EPI; LACI

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC217756 representing NM\_001177320

Red=Cloning site Blue=ORF Orange=Stop codon

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC}$ 

GCCGCGATCGCC

TTGGTTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001177320

**Insert Size:** 708 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## Tfpi (NM\_001177320) Mouse Untagged Clone - MC217756

**OTI Disclaimer:** 

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001177320.1</u>, <u>NP 001170791.1</u>

RefSeq Size: 2993 bp
RefSeq ORF: 708 bp
Locus ID: 21788
Cytogenetics: 2 D

**Gene Summary:** 

Inhibits factor X (X(a)) directly and, in a Xa-dependent way, inhibits VIIa/tissue factor activity, presumably by forming a quaternary Xa/LACI/VIIa/TF complex. It possesses an antithrombotic action and also the ability to associate with lipoproteins in plasma (By similarity).

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) differs in the 5' UTR, 3' UTR, and 3' coding sequence compared to variant 1. The resulting isoform (b) has a shorter and distinct C-terminus compared to isoform a. Variants 2 and 3 both encode isoform b. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.