

Product datasheet for MC218254

Morf4l1 (BC017619) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Morf4l1 (BC017619) Mouse Untagged Clone

Tag: Tag Free
Symbol: Morf4l1

Synonyms: TEG-189, MRG15, MORFRG15, mKIAA4002

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin
Fully Sequenced ORF: >BC017619

Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATTACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:Sgfl-MlulACCN:BC017619

Insert Size: 972 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Morf4l1 (BC017619) Mouse Untagged Clone - MC218254

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>BC017619</u>, <u>AAH17619</u>

RefSeq Size: 1246 bp
RefSeq ORF: 971 bp
Locus ID: 21761
Cytogenetics: 9 E3.1

Gene Summary: Component of the NuA4 histone acetyltransferase complex which is involved in

transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote

interaction of the modified histones with other proteins which positively regulate

transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the mSin3A complex which acts to repress transcription by deacetylation of nucleosomal histones. Required for homologous recombination repair (HRR) and resistance to mitomycin C (MMC).

Involved in the localization of PALB2, BRCA2 and RAD51, but not BRCA1, to DNA-damage foci

(By similarity).[UniProtKB/Swiss-Prot Function]