

Product datasheet for **MC218803**

Fnbp1 (NM_001038700) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Fnbp1 (NM_001038700) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Fnbp1
Synonyms:	1110057E06Rik; 2210010H06Rik; FBP1; Fbp17
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC218803 representing NM_001038700
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGAGCTGGGCACTGAGCTCTGGGATCAGTTTGACAACCTGGAAAAACATACACAGTGGGGAATCGATA
 TTCTTGAGAAATACATCAAGTTTGTCAAGGAGAGGACGGAGATTGAGCTCAGCTATGCCAAGCAACTCAG
 GAATCTTTCAAAGAAATACCAACCTAAGAAGAAGCTCGAAGGAAGAGGAGGAGTACAAGTACACGGCTTGC
 AAAGCCTTTCTTCCACCCTGAATGAGATGAATGACTACGCCGGGCAGCAGAGGTCATCTCTGAGAACA
 TGACGTCACAGATCACGGTGGACCTGATGCGCTACGTTCCAGGAGCTGAAGCAGGAGAGGAAATCGAACTT
 CCATGATGGACGGAAGGCTCAGCAGCACATAGAAACGTGTTGGAAGCAACTGGAGTCAAGTAAGAGGAGG
 TTTGAGCGGGACTGTAAGGAAGCCGACCGGGCACAGCAGTACTTCGAGAAAATGGACGCTGACATCAACG
 TGACCAAGCGGATGTGAAAAGGCACGACAACAAGCTCAGATACGCCAGCAAATGGCAGAGGACAGCAA
 AGCAGATTACTCCTTAATCCTGCAGAGATTCACCAGGAGCAATGGGAATACTACCATACCCACATCCCC
 AACATCTTCCAGAAAATACAAGAGATGGAGGAGAGGCGGATTGTGCGTATTGGGGAGTCCATGAAGACGT
 ACGCAGAGGTGGACCGGACGGTATACCCATCATCGGGAAGTGCCTGGACGGGATAGTGAAGCGGCCGA
 GTCTATCGACCAGAAAAACGACTCCCAGCTGGTCTAGAACCTATAAGTCAGGATTCGAGCCTCCTGGA
 GACATTGAATTCGAAGATTACACACAGCCAATGAAACGCACAGTGTGAGCAACAGCCTTCCAGCTCCA
 AAGAAGGCAAGCCTGAGCTCAGATTTGGCGCAAGTCCAGAGGCAAGCTCTGGCCATTATCAAGAAAAA
 CAAGGGTGTACACCAGAAGACTTCAGCAACTCCCACCTGAGCAGAGAAGGAAAAAACTACAACAGAAA
 GTTGACGATCTCAATAGAGAGATACAGAAGGAGACGGACCAGAGAGATGCCATCACCAAAATGAAAGATG
 TGTACCTAAAGAACCCTCAGATGGGAGACCCAGCCAGCCTGGACCAGAAGCTCACCGAAGTCACCCAGAA
 CATAGAGAAAACGCGCTGGAGGCTCAGAAGTTTGAGGCCTGGCTGGCTGAGGTAGAAGGCAGACTCCCA
 GCTCGGAGTGAGCAGGCACGCCGCGCAGAGTGGACTGTATGATGGCCAGACACACCAGACGGTCACTAACT
 GTGCCAGGACCGGGAGAGCCAGATGGTAGTTACACAGAGGAGCAAAGCCAGGAGAGCGAGCACAAAGT
 CCTGGCCCCGGATTTGACGATGAATTTGATGATGAGGAGCCGCTTCCAGCCATAGGGACCTGCAAGGCC
 CTCTACACATTTGAAGGTCAGAACGAAGGCACCATTTCCGTAGTTGAAGGAGAGACGCTGAGCGTGATTG
 AAGAGGACAAAGGCGATGGTGGACTCGCATCCGAGAAAATGAAGACGAGGAGGGTTACGTCCCCACTTC
 CTACGTCGAAGTCTATTTAGACAAAAACGCCAAAGGTTCC**TAA**

ACGCGTACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001038700

Insert Size: 1653 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001038700.2 , NP_001033789.1
RefSeq Size:	4545 bp
RefSeq ORF:	1653 bp
Locus ID:	14269
UniProt ID:	Q80TY0
Cytogenetics:	2 B
Gene Summary:	<p>Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during the late stage of clathrin-mediated endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also enhances actin polymerization via the recruitment of WASL/N-WASP, which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. May act as a link between RND2 signaling and regulation of the actin cytoskeleton. May be required for the lysosomal retention of FASLG/FASL (By similarity).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (1) lacks two alternate, in-frame exons, compared to variant 3. The resulting protein (isoform a) is shorter when it is compared to isoform c.</p>