

Product datasheet for **MC218816**

Psap (NM_001146123) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Psap (NM_001146123) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Psap
Synonyms:	AI037048; SGP; SGP-1
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >MC218816 representing NM_001146123
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGTACGCCCTCGCCCTCTTCGCCAGCCTTCTGGCCACCCTCTGACCAGCCCTGTCCAAGACCCGAAGA
 CATGCTCTGGGGCTCAGCAGTCTGTGCAGAGATGTGAAGACGGCGGTGGACTGTGGGGCCGTGAAGCA
 CTGCCAGCAGATGGTCTGGAGCAAGCCACAGCGAAATCCCTTCCTTGGACATATGCAAACTGTTGTC
 ACCGAAGCTGGAACTTCTGAAAGATAATGCTACGCAGGAGGAGATCCTTCATTACCTGGAGAAGACCT
 GTGAGTGGATTACGACTCCAGCCTGTCCGGCTCGTGCAAGGAGGTGGTTGACTCTTACCTGCCTGTCAT
 CCTGGACATGATTAAGGGCGAGATGAGCAACCCTGGGGAAGTGTGCTCTGCGCTCAACCTCTGCCAGTCC
 CTTCAGGAGTACTGGCCGAGCAAAACCAGAAACAGCTTGAGTCCAACAAGATCCCGGAGGTGGACATGG
 CCCGTGTGGTTGCCCCCTCATGTCCAACATCCCTCTCCTGCTGTACCCTCAGGATCACCCCGCAGCCA
 GCCCAACCTAAGGCTAACGAGGACGTCTGCCAGGACTGTATGAAGCTGGTGTCTGATGTCCAGACTGCT
 GTGAAGACCAACTCCAGCTTTATCCAGGGCTTCGTGGACCACGTGAAGGAGGATTGTGACCGCTTGGGGC
 CAGGCGTGTCTGACATATGCAAGAACTACGTGGACCAGTATCCGAGCAACCCAAGGAAATCTGTGTGCT
 GGCTGGCTTCTGTAATGAGGTCAAGAGAGTGCCAAATGAAGACTCTGGTCCCTGCCACCGAGACCATAAG
 AACATCCTCCCTGCCCTGGAGATGATGGACCCCTATGAGCAGAATCTGGTCCAGGCCACAATGTGATTT
 TATGCCAGACCTGTCAGTTTGTGATGAATAAGTTTTCTGAGCTGATTGTCAATAATGCCACTGAGGAGCT
 CCTAGTTAAAGGTTTGTGCAACGCATGCGCACTGCTCCCCGATCCTGCCAGAACCAAGTCCAGGAGGTG
 GTGGGAACATTTGGCCCTCCCTGTTGGACATCTTATCCATGAGGTAACCCAGCTCTGTGCGGTG
 TGACTGGCTCAAAGAGCCACACCGCCAAAGCAGCCCGCACAGCCCAAGCAGTTCGGCATTGCCCGCCAT
 GTGCCCTCAGAAAGTGGTGGTTCTGTGAGGTGTGCAAGAACTGGTCTCTATTTGGAACATAACC
 TGGAGAAAAACAGCACCAAGGAGGAAATCCTGGCCGCACTTGAGAAGGGCTGCAGCTTCTGCCAGACCC
 TTACCAGAAGCAGTGGATGACTTTGTGGCTGAGTATGAGCCCTTGTATTGGAGATCCTCGTGGAAAGT
 ATGGATCTGGATTTGTGTGCTCGAAAATTGGAGTTTGCCTTCTGCCTATAAGCTGCTGCTGGAAACCG
 AGAAGTGTGTCTGGGCCCTAGCTACTGGTGTGAGAACATGGAGACTGCCGCCGATGCAATGCTGTGCA
 TCATTGCAAACGCCATGTGTGGAAC**TAG**

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** Sgfl-Mlul
- ACCN:** NM_001146123
- Insert Size:** 1638 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_001146123.1](#), [NP_001139595.1](#)

RefSeq Size: 2640 bp

RefSeq ORF: 1638 bp

Locus ID: 19156

UniProt ID: [Q61207](#)

Cytogenetics: 10 30.02 cM

Gene Summary: This gene encodes a multifunctional glycoprotein that plays a role in the intracellular metabolism of various sphingolipids or secreted into the plasma, milk or cerebrospinal fluid. The encoded protein undergoes proteolytic processing to generate four different polypeptides known as saposin A, B, C or D, that are required for the hydrolysis of certain sphingolipids by lysosomal hydrolases. Alternately, the encoded protein is secreted into body fluids where it exhibits neurotrophic and myelinotrophic activities. A complete lack of the encoded protein is fatal to mice either at the neonatal stage or within the first month due to severe leukodystrophy and sphingolipid accumulation. Alternative splicing results in multiple transcript variants encoding different isoforms that may undergo similar processing to generate the mature saposins. [provided by RefSeq, Sep 2015]
Transcript Variant: This variant (5) lacks an exon in the coding region and uses an alternate in-frame splice site, compared to variant 2, resulting in a shorter protein (isoform E), compared to isoform B.