

## Product datasheet for **MC219698**

### **Fnbp1 (NM\_001177648) Mouse Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Fnbp1 (NM_001177648) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Fnbp1
Synonyms:	1110057E06Rik; 2210010H06Rik; FBP1; Fbp17
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

**Fully Sequenced ORF:** >MC219698 representing NM\_001177648  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGCC**

ATGAGCTGGGCACTGAGCTCTGGGATCAGTTTGACAACCTGGAAAAACATACACAGTGGGGAATCGATA  
 TTCTTGAGAAATACATCAAGTTTGTCAAGGAGAGGACGGAGATTGAGCTCAGCTATGCCAAGCAACTCAG  
 GAATCTTTCAAAGAAATACCAACCTAAGAAGAACTCGAAGGAAGAGGAGGAGTACAAGTACACGGCTTGC  
 AAAGCCTTTCTTCCACCCTGAATGAGATGAATGACTACGCCGGGCAGCAGAGGTCATCTCTGAGAACA  
 TGACGTCACAGATCACGGTGGACCTGATGCGCTACGTTCCAGGAGCTGAAGCAGGAGAGGAAATCGAACTT  
 CCATGATGGACGGAAAGGCTCAGCAGCACATAGAAACGTGTTGGAAGCAACTGGAGTCAAGTAAGAGGAGG  
 TTTGAGCGGGACTGTAAGGAAGCCGACCGGGCACAGCAGTACTTCGAGAAAATGGACGCTGACATCAACG  
 TGACCAAGCGGATGTGGAAGCCGACCGGACACAAGCTCAGATACGCCAGCAAATGGCAGAGGACAGCAA  
 AGCAGATTACTCCTTAATCCTGCAGAGATTCAACCAGGAGCAATGGGAATACTACCATACCCACATCCCC  
 AACATCTTCCAGAAAATACAAGAGATGGAGGAGAGGCGGATTGTGCGTATTGGGGAGTCCATGAAGACGT  
 ACGCAGAGGTGGACCGGACGGTATACCCATCATCGGGAAGTGCCTGGACGGGATAGTGAAGCGGCCGA  
 GTCTATCGACCAGAAAAACGACTCCCAGCTGGTCTAGAACCTATAAGTCAGGATTCGAGCCTCCTGGA  
 GACATTGAATTCGAAGATTACACACAGCCAATGAAACGCACAGTGTGAGCAACAGCCTTTCCAGCTCCA  
 AAGAAGGCAAGCTGAGCTCAGATTTGGCGGCAAGTCCAGAGGCAAGCTCTGGCCATTATCAAGAAAAA  
 CAAGCTTATGTCCTTTTAAATCCCCCATCAGCCTCCCCCTCCGCCCTGCCTGCCTCACCTCT  
 GCTGTTCCCAACGGCCCCAGTCTCCAAGCAGCCAAAGGAACCCCTCTCCACCGCTTCAACGAGTTCA  
 TGACTCCAAACCCAAAATCCACTGCTTCCGGAGCCTAAAGCGTGGGCTTCTCTCAAGCTGGGTGTCAC  
 ACCAGAAGACTTCAGCAACTTCCACCTGAGCAGAGAAGGAAAAAACTACAACAGAAAGTTGACGATCTC  
 AATAGAGAGATACAGAAGGAGACGGACCAGAGAGATGCCATACCAAAAATGAAAGATGTGTACCTAAAGA  
 ACCCTCAGATGGGAGACCCAGCCAGCCTGGACCAGAAGCTCACCGAAGTCACCCAGAACATAGAGAACT  
 GCGGCTGGAGGCTCAGAAGTTTGGGCTGGCTGGCTGAGGTAGAAGGCAGACTCCAGCTCGGAGTGAG  
 CAGGCACGCCGAGAGTGGACTGTATGATGGCCAGACACACCAGACGGTCACTAACTGTGCCAGGACC  
 GGGAGAGCCAGATGGTAGTTACACAGAGGAGCAAAGCCAGGAGAGCGAGCACAAGGTCCTGGCCCCGGA  
 TTTGACGATGAATTTGATGATGAGGAGCCGCTTCCAGCCATAGGGACCTGCAAGGCCCTACACATTT  
 GAAGGTCAGAACGAAGGCCACATTTCCGTAGTTGAAGGAGAGACGCTGAGCGTATTGAAGAGGACAAAG  
 GCGATGGGTGACTCGCATCCGCAGAAAATGAAGACGAGGAGGTTACGTCCCACTTCTACGTCGAAGT  
 CTATTTAGACAAAACGCCAAAGGTTCT**AA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-MluI

**ACCN:** NM\_001177648

**Insert Size:** 1851 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_001177648.1</a></u> , <u><a href="#">NP_001171119.1</a></u>
<b>RefSeq Size:</b>	4743 bp
<b>RefSeq ORF:</b>	1851 bp
<b>Locus ID:</b>	14269
<b>UniProt ID:</b>	<u><a href="#">Q80TY0</a></u>
<b>Cytogenetics:</b>	2 B
<b>Gene Summary:</b>	<p>Required to coordinate membrane tubulation with reorganization of the actin cytoskeleton during the late stage of clathrin-mediated endocytosis. Binds to lipids such as phosphatidylinositol 4,5-bisphosphate and phosphatidylserine and promotes membrane invagination and the formation of tubules. Also enhances actin polymerization via the recruitment of WASL/N-WASP, which in turn activates the Arp2/3 complex. Actin polymerization may promote the fission of membrane tubules to form endocytic vesicles. May act as a link between RND2 signaling and regulation of the actin cytoskeleton. May be required for the lysosomal retention of FASLG/FASL (By similarity).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (3) represents the longest transcript and it encodes the longest protein (isoform c).</p>