

Product datasheet for **MC220071**

Srpk1 (NM_016795) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Srpk1 (NM_016795) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Srpk1
Synonyms:	AU017960
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >MC220071 representing NM_016795
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGAGCGGAAAGTGCTCGCGCTCCAGGCCGAAAGAAAAGGACCAAGGCCAAGAAGGACAAAGCCAAA
 GGAAGCCTGAAACTCAGCACCGAGGCTCTGCACCCCACTCCGAGAGCGACATCCCAGAGCAGGAGGAGGA
 GATTCTAGGGTCTGACGATGACGAGCAGGAAGACCCCAATGACTACTGTAAGGAGGTTATCATCTTGTG
 AAAATTGGAGATCTATTTAACGGGAGATACCATGTGATTCGAAAAATTGGGCTGGGACACTTTTCCACAG
 TGTGGTTATCATGGGATATTCAGGGAAAGAAGTTGTAGCAATGAAAGTAGTTAAAAGTCTGAGCATTAA
 CACAGAAACAGCACTAGATGAAATCCGGCTGCTGAAATCGGTTCCCAACTCAGACCCGAATGATCCAAAT
 GGAGAAATGGTTGTTCACTACTAGATGACTTTAAAATATCAGGCGTTAATGGAACACATATCTGCATGG
 TGTTTGAGGTTTTGGGGCATCTACTCAAGTGGATCATCAAGTCCAATATCAGGGGCTCCGCTGCC
 TTGTGTCAAAAAAATTATTCAGCAAGTGCTACAGGGCCTGGATTACTTACACACCAAGTGCCGGATCATC
 CACACGGACATCAAAACCAGAGAACATCTTGTGTCTGTGAATGAGCAGTACATTCGAAGGCTGGCTGCAG
 AGGCCACAGAGTGGCAGCGCTCAGGAGCTCCCCACCATCGGGGTCTGCAGTCAGCACTGCTCCACAGCC
 TAAGCCAGCTGACAAAATGTCAAAGAATAAGAAGAAGAAATTGAAGAAGAAGCAGAAGCGCCAGGCAGAA
 TTAGTAGAGAAGCGAATGACAGGAAATGAGGAAATGGAGAAAGAGTCGGGCCCTGGGCAAAAAAGACCTA
 ACAAGCAAAGAAGATCAGAGAGTCTGTTGACAGACCCCTGACAGAGAACCCACCTAATAAAATGACCCA
 AGAGAACTTGAAGAGTCAAATTCATTGGCCAGGACCAGACACTCACGGAGCGTGGTGGAGAGGGTGGT
 GCGCCAGAGATTAATTGCAATGGAGTGAATGGAGTCGTTAATTACCCTGAGAACAGTAATAATGAGACAC
 TGAGGCATAAAGAGGATCTGCATAATGCTAATGACTGTGATGTCCACACTTTGAAGCAGGAACCTAGTTT
 CCTAAACTCTTCAAATGGAGACAGCAGTCCATCTCAAGACACAGACTCTTGACACCCACCGCCTCTGAG
 ACCATGGTGTGCCAATCCTCTGCAGAGCAGTCACTCACCCGACAGGACATCACCCAGCTGGAGGAGAGCA
 TTCGGGCAGATACACCTTCTGGGGATGAGCAGGAGCCCAATGGAGCCCTGGACAGCAAAGGAAAATTCTC
 TGCTGGAAATTTTCTTATTAATCCCCTTGAGCCAAAAATGCAGAGAAGCTCCAAGTGAAGATCGCAGAC
 CTCGGGAACGCCTGCTGGGTGCACAAGCATTCTACTGAAGATATCCAGACACGGCAGTATCGGTCTTTGG
 AAGTCTGATAGGATCTGGCTACAACACCCTGCTGACATCTGGAGCACAGCATGCATGGCCTTTGAACT
 AGCCACAGGGGACTATTTGTTTGGCCTATTGAGGGGAGGATTACACACGAGATGAAGACCACATCGCC
 CTGATCATAGAATCTGGGGAAGTGCCTCGCAAGCTCATTGTGCGAGGAAAATACTCCAAGGAATTTT
 TCACCAAAAAAGGTGACCTGAAGCACATACCAAGTTGAAGCCTTGGGGTCTTCTGGAGGTTCTGGTGGA
 GAAATATGAGTGGCTCAGGAGGAGGCTGCCGGCTTACAGATTTCTGTGCTGCCATGTTGGAGCTTATG
 CCTGAGAAGAGAGCCACTGCTGCTGAGTGTCTCCGGCATCCTTGCTAAACTCT**AA**

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_016795
- Insert Size:** 1947 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_016795.3](#), [NP_058075.2](#)

RefSeq Size: 2484 bp

RefSeq ORF: 1947 bp

Locus ID: 20815

UniProt ID: [O70551](#)

Cytogenetics: 17 A3.3

Gene Summary: Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine residues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phosphorylation of SR splicing factors and the regulation of splicing. Plays a central role in the regulatory network for splicing, controlling the intranuclear distribution of splicing factors in interphase cells and the reorganization of nuclear speckles during mitosis. Can influence additional steps of mRNA maturation, as well as other cellular activities, such as chromatin reorganization in somatic and sperm cells and cell cycle progression. Phosphorylates SFRS2, ZRSR2, LBR and PRM1. Phosphorylates SRSF1 using a directional (C-terminal to N-terminal) and a dual-track mechanism incorporating both processive phosphorylation (in which the kinase stays attached to the substrate after each round of phosphorylation) and distributive phosphorylation steps (in which the kinase and substrate dissociate after each phosphorylation event). The RS domain of SRSF1 binds first to a docking groove in the large lobe of the kinase domain of SRPK1. This induces certain structural changes in SRPK1 and/or RRM2 domain of SRSF1, allowing RRM2 to bind the kinase and initiate phosphorylation. The cycles continue for several phosphorylation steps in a processive manner (steps 1-8) until the last few phosphorylation steps (approximately steps 9-12). During that time, a mechanical stress induces the unfolding of the beta-4 motif in RRM2, which then docks at the docking groove of SRPK1. This also signals RRM2 to begin to dissociate, which facilitates SRSF1 dissociation after phosphorylation is completed. Can mediate hepatitis B virus (HBV) core protein phosphorylation. It plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgRNA) without affecting the formation of the viral core particles. Can induce splicing of exon 10 in MAPT/TAU (By similarity).[UniProtKB/Swiss-Prot Function]