

Product datasheet for **MC220075**

Tcf3 (NM_001164152) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Tcf3 (NM_001164152) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Tcf3
Synonyms:	A1; AA408400; ALF2; AW209082; bHLHb21; E2A; E12; E12/E47; E47; KA1; ME2; Pan1; Pan2; TCF-3; Tcfe2a
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >MC220075 representing NM_001164152
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGATGAACCACTCTCAGAGAATGGCACCCGTGGGCTCTGACAAGGAAGTGAAGTACCTCCTGGACTTCA
 GCATGATGTTCCCGCTACCTGTGGCCAATGGGAAGAGCCGGCCCGCTCCCTCGGGGAACCCAGTTTGC
 AGGCTCAGGACTGGAGGACCGACCCAGCTCAGGCTCCTGGGGCAGCAGTGACCAGAACAGTTCTTCTTT
 GACCTAGCCGGACATACAGCGAAGGTGCCACTTCAGTACTCCACAGCAGCTGCCGCTTCCACGT
 TCCTAGGAGCTGGGCTTGGAGGCAAGGGCAGTGAGCGAATGCCTATGCCACCTTTGGGAGAGACACCAG
 TGTTGGCACCTTGAGTCAGGCTGGCTTCTGCCAGGTGAGCTGAGCCTCAGCAGTCCCGGCCACTGTCC
 CCATCGGGCATCAAGAGCAGTCCAGTATTACCCCTCATTCCCAGCAACCCTCGTCGGAGAGCTGCAG
 ATGGTGGCCTGGCAGATACTCAGCCGAAGAAGGTCCGGAAGGTTCCGCCTGGTCTCCCTTCTCGGTGTA
 TCCGCCAGCTCAGGTGACAGCTACAGCAGGGATGCTGCAGCCTACCCCTCCGCCAAGACCCCCAGCAGC
 GCTTACCCCTCCCTTCTACGTGGCAGATGGCAGCCTGCACCATCAGCTGAGCTCTGGAGTACGCCTA
 GCCAGGTGGGCTTTGGGCCCAGTGTAGGTGACGGCTTCTCCCTCTGCCCTTGACCCGGGAGCAGCTC
 CGTGGGCACTGGTACCTTTGGGGCCCTCCAGCAGCAGGATCGCATGGGCTACCAGCTGCATGGATCTGAG
 GTTAATGGCTCGCTCCAGCTGTATCCAGCTTTTCGGCTGCCCTGGCACTTACAGTGGGACTTCCGGCC
 ACACGCCCCCTGTGAGTGGGGCCGACGTGAAAGCCTCCTAGGCACCCGAGGGACTACAGCCAGCAGCTC
 AGGGGATGCCCTTGGGAAGGCACTGGCCTCGATCTACTCCCGGATCACTCCAGCAATAATTTCTCACCT
 AGCCCCCAACGCCTGTGGGTTACCCAGGGCCTGCCAGGGACATCACAGTGGCCCCGGGAGGAGCGC
 CCAGTGCCTTATCCCCAACTACGATGCAGGTCTCCATGGCCTGAGCAAGATGGAGGACCGCTTGGACGA
 GGCCATCCATGTCTGCGAAGCCACGCTGTTGGCACCCTAGCGATCTCCATGGGCTTTTGCTTGGCCAT
 GGCGCACTGACCACGAGCTTACCGGCCCATGTCACTGGGCGGGCGCATGCCGCTGTGCGGGGAA
 GCCATCTGAGGAGGGCCTCACAAGTGGGGCCAGTCTTTTGCATAACCATGCCAGCCTCCCGAGCCAGCC
 CAGTTCCTCCTGACCTCTCACAGAGACCTCCCGACTCTATAGTGGACTCGGGAGGGCAGGCACAACA
 GCGGGTGCACGAGATCAAGCGGGAGGAGAAAGAGGATGAGGAAATCGCATCAGTAGCCGACGCCGAAG
 AGGACAAGAAGGACCTGAAGTCCCACGCACGCGCACCAGCAGTACAGATGAGGTGCTGTCCCTGGAGGA
 GAAGGACCTGAGGACCGGGAGAGGCGTATGGCCAATAACGCTCGGGAGCGGGTGCCTGTGCGGGACATT
 AACGAGGCCTTCCGGGAGCTGGGCCGATGTGCCAGCTGCACCTCAAGTCGGATAAGGCGCAGACCAAGC
 TGCTCATCTGCAGCAGCGGTGCAGGTATCCTGGGCCTGGAGCAGCAGGTGCGAGAACGCAACCTGAA
 CCCCAAAGCAGCTGCTTGAAGCGGAGGGAGGAGGAGAAGGTGTCTGGCGTGGTGGGGACCCACAGCTG
 GCCCTGTGAGCCGCCACCCGGGCTGGGTGAGGCCACAACCCAGCCGGGCACCTG**TGA**

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI

ACCN: NM_001164152

Insert Size: 1950 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001164152.1, NP_001157624.1</u>
RefSeq Size:	3308 bp
RefSeq ORF:	1950 bp
Locus ID:	21423
Cytogenetics:	10 39.72 cM
Gene Summary:	Transcriptional regulator. Involved in the initiation of neuronal differentiation. Heterodimers between TCF3 and tissue-specific basic helix-loop-helix (bHLH) proteins play major roles in determining tissue-specific cell fate during embryogenesis, like muscle or early B-cell differentiation. Dimers bind DNA on E-box motifs: 5'-CANNTG-3'. Binds to the kappa-E2 site in the kappa immunoglobulin gene enhancer. Binds to IEB1 and IEB2, which are short DNA sequences in the insulin gene transcription control region.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (6) has an alternate splice site and an alternate penultimate coding exon, compared to variant 1. The resulting isoform (6) lacks an internal aa and has a distinct segment in the C-terminal region, compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.