

Product datasheet for **MC220165**

Pip5k1c (NM_008844) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pip5k1c (NM_008844) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Pip5k1c
Synonyms:	A1115456; A1835305; Pip5klgamma
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NM_008844.2
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGAGCTAGAGGTGCCGACGAGGCGGAGAGCGCCGAAGCGGGGCGGTGACGGCGGAAGCGGCCTGGT
 CTGCGGAGAGTGGGGCGGCGCAGGTATGACCCAGAAGAAGGCTGGCCTTGCAGAGGCCCTTGGTGAC
 AGGGCAGCCAGGCCCTGGCCATGGGAAGAAGCTGGGCCACCGAGGCGTGATGCGTCGGGAGAGACTACG
 TATAAGAAGACCACCTCGTCCACCTGAAGGGCGCCATCCAGCTGGGGATCGGGTACACGGTGGCAACC
 TGAGCTCTAAGCCGAGCGGGACGTGCTCATGCAGGACTTCTACGTGGTGGAGAGCATCTTCTCCCCAG
 TGAAGGGAGCAACCTCACCCCTGCCACCATTCCAGGATTTCCGCTTCAAGACCTATGCACCTGTTGCC
 TTCGCTACTTCCGGGAGCTCTTGGCATCCGTCCAGATGACTATTTGACTCCCTGTGCAATGAGCCAC
 TCATCGAGCTCTCAACCCGGGGCCAGCGGCTGTCTTCTACGTACCAGCGACGACGAGTTCATCAT
 CAAGACTGTCATGCACAAGGAGGACAGATTCTGCAGAAGCTGCTGCCTGGCTACTACATGAATCTCAAC
 CAGAACCACGGACGCTGCTGCCAAGTTCTATGGGCTGTACTGCGTGCAGTCTGGTGGCAAGAATCC
 GCGTGGTGGTCATGAACAATGTGCTGCCCGTGTGCTCAAAATGCACCTTAAGTTCGACCTCAAGGGCTC
 CACGTACAAGCGCAGGGCCAGCAAGAAGGAGAAGGAGAAGAGCCTGCCACCTACAAGGACCTGGACTTC
 ATGCAGGACATGCCCGAGGGGCTGCTGCTGGACTCCGACACCTTTGGCGCCCTGGTCAAGACGCTGCAGC
 GAGACTGCCTGGTGTGGAGAGCTTCAAGATAATGGACTACAGCCTGCTGCTGGGCGTGCACAACATCGA
 TCAGCAGGAGCGAGAGCGCCAGGCCGAGGGCGCCAGAGCAAGGGCGATGAGAAGCGGCCCTGGCCAG
 AAGGCTGTGATCCACGGCCATGGAGTCTATCCAGGGCGGAGCTGCCCGTGGGGAGGCCATTGAGACAG
 ATGACACGATGGGTGGGATTCCAGCAGTGAATGGGCGGGGAGCGACTGCTCTCCACATCGGGATCAT
 TGATATTCTGCAGTCTACAGTTTCATCAAGAAGTTAGAACACACCTGGAAGGCCCTCGCCATGATGGG
 GACACTGTCTCAGTCCACCGGCCAGTCTTATGCAGAGCGCTTCTTCAAGTTCATGAGCAGCACGGTGT
 TCCGGAAGAGTTCCTCCCTGAAGTCTCTCCATCCAAGAAAGGACGTGGTGCCTGCTGGCGGTCAAACC
 CCTGGGGCCACTGCTGCCTTCTCAGCCAGCCAGATCCCAGCGAGAGAGAGGATGTGCAGTATGACCTG
 CGGGGGCCCGCAGTACCCACGCTTGAGGATGAAGGCCGGCTGACCTCCTGCCCTGCACCCACCGT
 CCTTTGAGGAAGCCACCACGCCTCCATCGCCACCACCCTGTGCTCCACCTCCCTCTCCATCCCAGAGCG
 GTCCTTTCAGATACATCGGAGCAGCCCGGTACAGGCGGCGCACGCAGTCTCAGGCCAGGATGGCCGG
 CCCAGGAGGAGCCCATGCGGAAGACCTGCAGAAGATAACTGTGCAGGTGGAGCCAGTGTGCGGTGTGG
 GGGTTGTCCCAAGGAGGAGGGTGCAGGAGTGGAGGTCCCCCATGTGGGGCATCGGCTGCAGCCTCTGT
 GGAATAGACGCTGCCAGCCAGGCTCAGAGCCTGCCAGCCAGGCTCAGATGAGGAGGATGCACCTCT
 ACAGACATCTATTTCCACCGACGAGAGGAGCTGGGTGTACTCCCCGCTTCACTATAGCGCGGGCCCG
 CCTCCGACGGCGAGAGCGACACATAA

ACGCGTACGCGGGCCGCTCGAGCAGAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAAGTTTAA

Chromatograms: https://cdn.origene.com/chromatograms/ja1495_f02.zip

Restriction Sites: SgfI-MluI

ACCN: NM_008844

Insert Size: 1986 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_008844.2</u> , <u>NP_032870.2</u>
RefSeq Size:	4287 bp
RefSeq ORF:	1986 bp
Locus ID:	18717
UniProt ID:	<u>O70161</u>
Cytogenetics:	10 C1

Gene Summary:

Catalyzes the phosphorylation of phosphatidylinositol 4-phosphate (PtdIns4P) to form phosphatidylinositol 4,5-bisphosphate (PtdIns(4,5)P₂). PtdIns(4,5)P₂ is involved in a variety of cellular processes and is the substrate to form phosphatidylinositol 3,4,5-trisphosphate (PtdIns(3,4,5)P₃), another second messenger. The majority of PtdIns(4,5)P₂ is thought to occur via type I phosphatidylinositol 4-phosphate 5-kinases given the abundance of PtdIns4P. Participates in a variety of cellular processes such as vesicle mediated transport, cell adhesion, cell polarization and cell migration. Together with PIP5K1A is required for phagocytosis, but they regulate different types of actin remodeling at sequential steps. Promotes particle attachment by generating the pool of PtdIns(4,5)P₂ that induces controlled actin depolymerization to facilitate Fc-gamma-R clustering. Mediates RAC1-dependent reorganization of actin filaments. Required for synaptic vesicle transport. Controls the plasma membrane pool of PtdIns(4,5)P₂ implicated in synaptic vesicle endocytosis and exocytosis. Plays a role in endocytosis mediated by clathrin and AP-2 (adaptor protein complex 2). Required for clathrin-coated pits assembly at the synapse. Participates in cell junction assembly. Modulates adherens junctions formation by facilitating CDH1 trafficking. Required for focal adhesion dynamics. Modulates the targeting of talins (TLN1 and TLN2) to the plasma membrane and their efficient assembly into focal adhesions. Regulates the interaction between talins (TLN1 and TLN2) and beta-integrins. Required for uropodium formation and retraction of the cell rear during directed migration. Has a role in growth factor- stimulated directional cell migration and adhesion. Required for talin assembly into nascent adhesions forming at the leading edge toward the direction of the growth factor. Negative regulator of T-cell activation and adhesion. Negatively regulates integrin alpha-L/beta-2 (LFA-1) polarization and adhesion induced by T-cell receptor. Together with PIP5K1A has a role during embryogenesis and together with PIP5K1B may have a role immediately after birth. [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) lacks an alternate in-frame exon in the 3' coding region, compared to variant 3. The encoded protein (isoform 1) is shorter, compared to isoform 3.

Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.