

Product datasheet for **MC220686**

Ddr1 (BC065998) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ddr1 (BC065998) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ddr1
Synonyms:	6030432F18, CD167a, PTK3A
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin



[View online »](#)

Fully Sequenced ORF: >BC065998
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGCATCGCC**

ATGGGGACAGGGACCCTCTCATCTCTACTGCTGCTACTCTTGGTGACAATTGGAGATGCTGACATGA
 AGGGACATTTTGACCCTGCCAAGTGCCGCTATGCCCTGGGCATGCAGGACCGCACCATTCTGACAGCGA
 TATCTCTGTGTCCAGCTCCTGGTCGACTCTACCCTGCCCGCCACAGCAGGCTGAAAGCAGTGATGGA
 GATGGGGCTTGGTGCCCTGCAGGGCTGTGTTCCCAAAGAAGAGGAGTACTTGCAGGTGGACCTTCGTA
 GGCTACACCTGGTGGCTCTGGTGGGCACCCAGGGCCGCATGCTGGGGTCTGGGCAAAGAGTTCTCCCG
 AAGCTATCGTTGCGTTACTCCCGAGATGGCCGCGCTGGATGGACTGGAAGGACCGCTGGGACAGGAG
 GTGATTCGGGTAACGAGGATCCCGGGGAGTAGTCTGAAGGACCTGGGCCCCCATGGTGGCCCGGC
 TGGTCCGCTTCTACCCAGGGCTGACCGGGTATGAGTGTCTGTCTTCGGGTGGAGCTCTATGGTGCCT
 CTGGCGGATGGACTCCTGTATACAGCCCCGCTGGGGCAGACCATGCAGTTATCTGAGGTGATGGTA
 CATCTCAATGATTCCACTTACGATGGATATACTGCTGGAGGGCTGCAATATGGCGGTCTGGGCCAGCTGG
 CAGATGGCGTGGTGGGCCTGGATGATTTAGGCAGAGCCAGGAGCTGCGGGCTGGCCAGGCTATGACTA
 TGTGGGATGGAGCAATCAGAGCTTCCCCACGGCTACGTGGAGATGGAGTTTGGTTGATCGGTTGAGG
 ACCTTCCAGACCATGCAGGTCCACTGTAACAACATGCACACTCTGGGAGCCCGCTACCAGGCGGGGTGG
 AATGCCGGTTTAAAAGGGTCCCGCCATGGCCTGGGAAGGAGAGCCTGTCCGCCATGCTCTGGGAGGCAG
 CCTTGGAGACCCAGAGCCCGGGCCATCTCAGTCCCCGGTGGCCACGTGGGCCGCTTCTGCAGTGC
 AGATTCCTCTTTGAGGTCCTTGGTACTCTTTCAGTGAATCTCTTTCATCTCAGATGTGGTGAACGACT
 CCTCTGACACCTTCCCACAGCCCCCTGGTGGCCACCTGGCCCGCTCCCACTTCCAGCAGCTTGGTA
 GCTGGAGCCCCGGGTCAACAGCCAGTGGCCAAGGCGGAGGGGAGCCCAACTGCCATCCTATTGGCTGC
 CTGGTGGCCATCATCCTGCTGCTGCTTCTCATATCGCGCTGATGCTCTGGAGGCTGCACTGGCGCCGGC
 TGCTCAGCAAGGCTGAGCGCCGCTGTTGGAGGAGGAGCTGACGGTTACCTTTCTGTCCCTGGGACAC
 CATCCTCATCAACAACCGCCAGGACCCCGAGAGCCACCCCTTACCAGGAGCCCGGCCTCGGGGACT
 CCACCCATTCTGCACCTGCGTCCCCAACGGCTCTGCCTGCAGTGGGACTATATGGAGCCCGAGAAGC
 CGGGTCCCGCTTCTACCCACCTCCCAGAACAGCGTCCCCATTATGCCGAAGCTGACATTGTCAC
 CCTGCAGGGCGTCACTGGGGCAACACCTACGCTGTGCCTGCACTGCCCCAGGGCGGTGGGGATGGG
 CCCCCAGAGTGGATTTCCCTCGGTACGGCTCCGCTTCAAGGAGAAGCTTGGCAGGGCCAATTTGGGG
 AGGTACACCTGTGTGAAGTAGAGGACCCCGCAGCCCTTCCCAATGGGAGCCATCCAGGGGAAGCTGGAC
 TCGAAAACCAGAGAAGCCTATGGCACCGCACCCACCGTCTCCATACTTGCCATTCCCAAGAGGCAAGT
 TCTCTGCGGGTGGGCTGGGCCTGCCGAGGAGCAGATACACACTCCCTTCCGTTTCTTCTTCCATCT
 GCCAAGTGCCACCCTCCAGCTGGCCCTGTGGTTGGGATCCTGCCTGACTTCTCCAGCTTCTTCTTGT
 CACACACTGGACGCCCGCAGCTGAGAATTAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCTGAAGAGGATCTGGCAGCAATGATATCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-MluI
ACCN: BC065998
Insert Size: 2061 bp
OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>BC065998</u> , <u>AAH65998</u>
RefSeq Size:	2582 bp
RefSeq ORF:	2060 bp
Locus ID:	12305
Cytogenetics:	17 18.7 cM
Gene Summary:	Tyrosine kinase that functions as cell surface receptor for fibrillar collagen and regulates cell attachment to the extracellular matrix, remodeling of the extracellular matrix, cell migration, differentiation, survival and cell proliferation. Collagen binding triggers a signaling pathway that involves SRC and leads to the activation of MAP kinases. Regulates remodeling of the extracellular matrix by up-regulation of the matrix metalloproteinases MMP2, MMP7 and MMP9, and thereby facilitates cell migration and wound healing, but also tumor cell invasion. Promotes smooth muscle cell migration, and thereby contributes to arterial wound healing. Phosphorylates PTPN11 (By similarity). Required for normal blastocyst implantation during pregnancy, for normal mammary gland differentiation and normal lactation. Required for normal ear morphology and normal hearing.[UniProtKB/Swiss-Prot Function]