

Product datasheet for **MC220818**

Kirrel3 (NM_001190912) Mouse Untagged Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | Expression Plasmids |
| Product Name: | Kirrel3 (NM_001190912) Mouse Untagged Clone |
| Tag: | Tag Free |
| Symbol: | Kirrel3 |
| Synonyms: | 1500010O20Rik; 2900036G11Rik; mKIAA1867; NEPH2; SST4 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |



[View online »](#)

| | |
|-------------------------------|--|
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | <u>NM_001190912.1</u> , <u>NP_001177841.1</u> |
| RefSeq Size: | 3734 bp |
| RefSeq ORF: | 2334 bp |
| Locus ID: | 67703 |
| UniProt ID: | <u>Q8BR86</u> |
| Cytogenetics: | 9 A4 |
| Gene Summary: | <p>Synaptic adhesion molecule required for the formation of target-specific synapses (PubMed:23637329, PubMed:26575286). Required for formation of target-specific synapses at hippocampal mossy fiber synapses. Required for formation of mossy fiber filopodia, the synaptic structures connecting dentate granule and GABA neurons. Probably acts as a homophilic adhesion molecule that promotes trans-cellular interactions and stabilize mossy fiber filopodia contact and subsequent synapse formation (PubMed:26575286). Required for the coalescence of vomeronasal sensory neuron axons (PubMed:23637329). May be involved in the hematopoietic supportive capacity of stroma cells; the secreted extracellular domain is directly responsible for supporting hematopoietic stem cells (PubMed:12665856). [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (C) lacks an in-frame exon in the 5' coding region and uses an alternate in-frame splice site in the 3' coding region, compared to variant A. The resulting protein (isoform C) is shorter than isoform A. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.</p> |