

## Product datasheet for **MC221009**

### **Ikbke (NM\_019777) Mouse Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Ikbke (NM_019777) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ikbke
Synonyms:	AW558201; IKK-E; IKK-i; IKKepsilon; Ikki
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**Fully Sequenced ORF:** >MC221009 representing NM\_019777  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGC**

ATGCAGAGTACCACTAACTACCTGTGGCATACTGATGACCTGCTAGGGCAGGGGGCCACTGCCAGTGTGT  
 ACAAGGCCCGAAACAAGAAATCCGGGGAGGTGGTTGCTGTAAAGGTCTTCAACTCAGCCAGCTATCGGCG  
 ACCTCCTGAGGTTGAGGTGAGGGAGTTTGGAGTCTGCGGAGGCTGAATCACCAGAACATCGTGAAGCTA  
 TTCGAGTGGAGGAAACGGGAGGAGCCGGCAGAGGTGCTAATCATGGAGTACTGCTCCAGTGGGAGCC  
 TGCTGAGCGTGTGGAAGACCTGAGAACAGTTCGGGCTTTCTGAAGAGGAGTTCCTAGTGGTGTGCG  
 CTGTGTGGTGGCTGGCATGAACACCTGCGGGAGAATGGCATTGTCCATCGGGACATCAAACCTGGGAAC  
 ATCATGCGCTGGTGGGCGAGGAGGGGAGAGCATCTATAAGCTGTCTGACTTCGGGGCTGCCCGCAAGC  
 TGGACGATGATGAGAAGTTTGTCTGTCTATGGTACAGAGGAATACCTGCACCCTGACATGTATGAGCG  
 TGCAGTGTGCGAAACCCAGCAAAAGGCATTTGGTGTGACTGTGGATCTCTGGAGTATTGGGGTGACC  
 CTGTACCACGCAGCCACAGGCAGTCTGCCCTTCATCCCCTTCGGTGGGCCCGGCCCAACAAAGAGATCA  
 TGTACAGAATCACCACAGAGAAGCCAGCCGGGGCCATTTAGGGACTCAGAAGCAGGAAAATGGTCCCTT  
 GGAGTGGAGCTACAGCCTCCCCATCACCTGTAGACTGTCCATGGGACTGCAGAACCAGCTGGTGGCCATC  
 CTGGCCAAACATCTGGAGGTGGAAGAGGATAAGTGCTGGGGCTTTGATCAGTTCTTCGCGGAGACCAGTG  
 ACATTCTGCAGCGAACGGTCATCCACGTCTTTTCCCTACCCAGGCCGTTTGCATCATGTCTACATCCA  
 CGCCACAACACGATTGCCATCTTTTGGAGGCTGTATATGAGCAGACCAACGTGACCCCAACACCAG  
 GAGTACCTCTTCGAGGGTCACCTTGTGTCCTTGAGCCAAGCCTCTCAGCCAGCACATCGCCACACAG  
 CTGCCAGCAGCCCTCTAACTCTGTTCAAGATGTCCAGCGACACACCTAAGGGGCTGGCCTTCAGGGACCC  
 TGCTCTGGATGTCCCAAAGTTTCGTCCCTAAGGTTGACCTACAGGCCGATTACAGCACAGCTAAGGGGGTG  
 CTGGGCGCTGGCTACCAGGCCCTGTGGCTGGCGGGGTCTGCTGGATGGACAGGCGTTGATGCTTCGGG  
 GGTTACATTGGGTCTGGAGGTGCTTCAGGACACGTGCCAGCAGACACTGGAGGTACACGGACAGCCCT  
 CCTCTACCTCAGCAGCAGCTGGGCACTGAAAGGTTGAGCAGTGGAGCGGGGATGCCTGACGTCCAGGAA  
 CGAAAGGAGGCCACAGAGCTAAGAACCAGGCTGCAGACTCTCTCAGAGATCCTGTCTAAATGTTCCACA  
 ATGTCACAGAAACCCAAAGGAGCCTGAGCTGTCTGGGTGAAGAGCTTTTAAAGAACGGGACCAGATTCA  
 TGAGGATAACAAAAGTATCCAGAAGATTCAGTGTGTTTGGACAAGATGCACTTCATCTACAAACAGTTC  
 AAGAAATCCAGGATGAGGCCAGGCTCAGCTACAATGAGGAGCAGATCCACAAGCTGGATAAGGTAATT  
 TCAGTCATCTAGCCAAGAGGCTGCTGCAGGTGTTCCAGGAGGAGTGTGTGCAGACGTATCAGGTGTGCT  
 GGTACACACGGCAAGCGGATGAGGCAGGTGCAGAGGGCCAGAACACCTGCATCTCATTGGCCACTCT  
 GTGGCCACCTGTAACCTCGGAAGCCCGGGAGGCCAGGAGAGTCTGAACAAGATCTTTGATCAGCTCCTTC  
 TGGACAGAGCTTCCGAACAGGGAGCTGAGGTGTACCGCAACCTATGGCTCCTCATCCCGGCCCTGATCC  
 GAAGGACCTGGTCTTCACATGCAGGAGCTTTGTAATGATATGAAGCTATTGGCCTTTGATCTCCAGGAC  
 AACAACCGACTCATCGAACGTTACATAGAGTTCCATCGGCACCAGATGTC**TGA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-MluI  
**ACCN:** NM\_019777  
**Insert Size:** 2154 bp

**OTI Disclaimer:** Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at [custsupport@origene.com](mailto:custsupport@origene.com) or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** [NM\\_019777.3](#), [NP\\_062751.2](#)

**RefSeq Size:** 3203 bp

**RefSeq ORF:** 2154 bp

**Locus ID:** 56489

**UniProt ID:** [Q9R0T8](#)

**Cytogenetics:** 1 E4

**Gene Summary:**

Serine/threonine kinase that plays an essential role in regulating inflammatory responses to viral infection, through the activation of the type I IFN, NF-kappa-B and STAT signaling. Also involved in TNFA and inflammatory cytokines, like Interleukin-1, signaling. Following activation of viral RNA sensors, such as RIG-I-like receptors, associates with DDX3X and phosphorylates interferon regulatory factors (IRFs), IRF3 and IRF7, as well as DDX3X. This activity allows subsequent homodimerization and nuclear translocation of the IRF3 leading to transcriptional activation of pro-inflammatory and antiviral genes including IFNB. In order to establish such an antiviral state, IKBKE forms several different complexes whose composition depends on the type of cell and cellular stimuli. Thus, several scaffolding molecules including IPS1/MAVS, TANK, AZI2/NAP1 or TBKBP1/SINTBAD can be recruited to the IKBKE-containing-complexes. Activated by polyubiquitination in response to TNFA and interleukin-1, regulates the NF-kappa-B signaling pathway through, at least, the phosphorylation of CYLD. Phosphorylates inhibitors of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. In addition, is also required for the induction of a subset of ISGs which displays antiviral activity, may be through the phosphorylation of STAT1 at 'Ser-708'. Phosphorylation of STAT1 at 'Ser-708' seems also to promote the assembly and DNA binding of ISGF3 (STAT1:STAT2:IRF9) complexes compared to GAF (STAT1:STAT1) complexes, in this way regulating the balance between type I and type II IFN responses. Protects cells against DNA damage-induced cell death. Also plays an important role in energy balance regulation by sustaining a state of chronic, low-grade inflammation in obesity, wich leads to a negative impact on insulin sensitivity. Phosphorylates AKT1. [UniProtKB/Swiss-Prot Function]