

## Product datasheet for MC223176

### Ptk2b (NM\_001162365) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Ptk2b (NM_001162365) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ptk2b
Synonyms:	CADTK; CAKB; CAKbeta; E430023O05Rik; FADK2; FAK2; PYK2; Raftk
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC223176 representing NM_001162365 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCCCGCATCGCC

ATGTCCGGGGTGTCTGAGCCCTTGAGCCGTGTAAAAGTGGGCACTTTACGCCGCCTGAGGGCCCCCAG  
AGCCCATGGTGGTGTACCAGTGGATGTGGAGAAGGAAGACGTGCGCATCCTCAAGGTCTGCTTCTACAG  
CAACAGCTTCAACCCAGGAAGAAGTCAAGCTTGTCAAATGCACAGTGCAGACAGAGATCCAGGAGATC  
ATCACCTCCATCCTCCTGAGTGGGCGAATAGGGCCCAACATCCAGCTGGTGAATGCTATGGGCTGAGGC  
TGAAGCAGATGAAGTCAGACGAGATCCACTGGCTGCACCCACAGATGACCGTGGGCGAAGTGCAGGACAA  
GTATGAATGTCTACACGTGGAAGCTGAGTGGAGGTATGACCTTCAAATCCGCTACTTGCCGGAAGACTTC  
ATGGAGAGCCTGAAAGAAGACAGGACCACATTGCTGTACTTTTATCAACAGCTCCGGAATGACTACATGC  
AACGCTACGCCAGCAAGGTCAAGTGAAGGCATGGCTCTGCAGCTGGGCTGTCTGGAGCTCAGGAGATTCTT  
CAAGGACATGCCCCACAATGCACTGGACAAAAAGTCCAACCTTTGAACTCCTGGAAAAAGAAGTCGGTCTG  
GACCTGTTTTTCCCAAAGCAGATGCAGGAAAACTTAAAGCCCAAGCAGTTCGGAAAGATGATCCAGCAGA  
CCTTCCAGCAGTATGCATCACTCCGGGAGGAAGAGTGTGTATGAAATCTTCAATACCTAGCGGGCTT  
TGCCAACATTGACCAGGAGACCTACCGCTGCGAACTCATTCAAGGATGGAACATTACTGTGGACCTGGTC  
ATCGGCCCTAAAGGCATCCGTGAGTCAAGTCAAGATACAAAGCCACCTGGCCGAGTTTAAAGC  
AGATCAAATCCATCAGGTGCCTCCATTGGAAGAGACCCAGGCAGTCTGCAGCTGGGCATCGAGGGTGC  
CCCCAGTCCCTGTCTATCAAAACGTCGCTCCCTGGCAGAGGCTGAGAACATGGCTGACCTCATAGATGGC  
TACTGCAGGCTGCAAGGAGAACATAAGGGCTCTCTCATCATGCATGCCAAGAAAGATGGTGAAGAAGGA  
ACAGCCTGCCTCAGATCCCCACACTAAACCTGGAGGCTCGCGGTGCGACCTCTCAGAAAGCTGCAGCAT  
AGAGTCAGACATCTATGCGGAGATTCCCGATGAGACCTGCGAAGACCAGGAGGTCCACAGTACGGTGT  
GCCCGTGAAGAAGTAGTTCTTAACCGCATTCTGGGTGAAGGCTTCTTTGGGGAGGCTATGAAGGTGTCT  
ACACGAACCACAAAGGGGAAAAAATTAATGTGGCCGTCAAGACCTGTAAGAAAGACTGTACCCAGGACAA  
CAAGGAGAAGTTCATGAGTGAGGCAGTGATCATGAAGAATCTTGACCACCTCACATCGTGAAGCTGATT



[View online »](#)

GGCATCATTGAAGAGGAACCCACCTGGATTATCATGGAAGTGTATCCTTATGGGGAGCTGGGACACTACC  
 TGAACGAAATAAAAACCTCCCTGAAGGTACCCACTCTGGTCTGTACACCCTACAGATATGCAAAGCCAT  
 GGCCTATCTGGAGAGCATCAACTGTGTGCACAGGGATATTGCTGTCCGGAACATCCTGGTGGCCTCTCT  
 GAGTGTGTGAAGCTGGGGACTTTGGGCTCTCCCGGTACATTGAGGACGAAGACTATTACAAAGCCTCTG  
 TGACCCGTCTACCCATCAAATGGATGTCCCCGAGTCCATCAACTCCGCCGCTTACAACCGCCAGTGA  
 TGCTGGATGTTTGTGTATGCATGTGGGAGATCCTCAGCTTTGGGAAGCAGCCTTTCTTCTGGCTCGAA  
 AATAAGGATGTCATCGGAGTGTGGAGAAAGGGACAGGCTGCCAAGCCCGAAGCTGTCCGCTGTCC  
 TTTACACACTCATGACTCGTGTCTGGGACTACGACCCAGTGACCGGCCCGCTTACGGAGCTTGTGTG  
 CAGCCTCAGTGACATTTATCAGATGGAGAAGGACATTGCCATAGAGCAAGAAAGGAATGCTCGTACCGA  
 CCCCCAAAATATTGGAGCCTACTACCTTTAGGAACCCCAACCCAGCCGCGCCCAAGTACAGAC  
 CTCCTCCACAGACCAACCTGCTGGCTCCTAAGCTGCAGTTCAGGTCCCTGAGGGTCTGTGTGCCAGCTC  
 TCCTACGCTTACCAGCCCTATGGAGTATCCATCTCCAGTTAACTCGTACACACCCACCTCTCCACCGG  
 CACAATGTCTTCAAGCGCCACAGCATGCGGGAGGAGGACTTATCCGGCCAGTAGCCGAGAAGAGGCC  
 AGCAGCTCTGGGAGGCAGAGAAGATCAAGATGAAGCAGGTCTAGAAAGACAGCAGAAGCAGATGGTGGA  
 AGATCCAGTGGCTGAGGCGAGAGGAAAGATGCTTGGACCCTATGGTTTATATGAATGACAAGTCCCCA  
 CTGACTCCAGAGAAGGAGCCGGCTACAGGCCCCACAGAAACCCCTCGCTCGGTGCACAGTCCATTC  
 AGCCACAGCCAACCTGGACAGGACCGATGACCTCGTGTACCACAATGTCATGACCTGGTGGAGGCTGT  
 GCTGGAAGTCAAGAAAGCTTGGCCAGTTGCCCCCTGAGGACTATGTGGTGGTGGTGAAGAACGTGGGG  
 CTGAACCTGCGGAAGCTCATCGGCAGTGTGGACGATCTTTGCCCTCCTTCCCGGCATCTTCGAGGACAG  
 AGATTGAAGGGACCCAGAAACTGCTCAACAAAGACCTGGCAGAGCTCATCAACAAGATGAAGTTGGCTCA  
 GCAGAACGCCGTGACGTCCCTGAGTGAGGACTGCAAGCGGCAGATGCTCACAGCGTCCCATACCTGGCT  
 GTGGATGCCAAGAACCTGCTGGATGCTGTGGACCAAGCAAGGTTGTGGCTAATCTGGCCACCCCGCCTG  
 CAGAGTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:**

Sgfl-Mlul

**ACCN:**

NM\_001162365

**Insert Size:**

3018 bp

**OTI Disclaimer:**

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:**

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:**

NM\_001162365.1, NP\_001155837.1

**RefSeq Size:**

4039 bp

**RefSeq ORF:**

3018 bp

Locus ID: 19229

Cytogenetics: 14 34.36 cM

**Gene Summary:** Non-receptor protein-tyrosine kinase that regulates reorganization of the actin cytoskeleton, cell polarization, cell migration, adhesion, spreading and bone remodeling. Plays a role in the regulation of the humoral immune response, and is required for normal levels of marginal B-cells in the spleen and normal migration of splenic B-cells. Required for normal macrophage polarization and migration towards sites of inflammation. Regulates cytoskeleton rearrangement and cell spreading in T-cells, and contributes to the regulation of T-cell responses. Promotes osteoclastic bone resorption; this requires both PTK2B/PYK2 and SRC. May inhibit differentiation and activity of osteoprogenitor cells. Functions in signaling downstream of integrin and collagen receptors, immune receptors, G-protein coupled receptors (GPCR), cytokine, chemokine and growth factor receptors, and mediates responses to cellular stress. Forms multisubunit signaling complexes with SRC and SRC family members upon activation; this leads to the phosphorylation of additional tyrosine residues, creating binding sites for scaffold proteins, effectors and substrates. Regulates numerous signaling pathways. Promotes activation of phosphatidylinositol 3-kinase and of the AKT1 signaling cascade. Promotes activation of NOS3. Regulates production of the cellular messenger cGMP. Promotes activation of the MAP kinase signaling cascade, including activation of MAPK1/ERK2, MAPK3/ERK1 and MAPK8/JNK1. Promotes activation of Rho family GTPases, such as RHOA and RAC1. Recruits the ubiquitin ligase MDM2 to P53/TP53 in the nucleus, and thereby regulates P53/TP53 activity, P53/TP53 ubiquitination and proteasomal degradation. Acts as a scaffold, binding to both PDPK1 and SRC, thereby allowing SRC to phosphorylate PDPK1 at 'Tyr-9', 'Tyr-373', and 'Tyr-376' (By similarity). Promotes phosphorylation of NMDA receptors by SRC family members, and thereby contributes to the regulation of NMDA receptor ion channel activity and intracellular Ca(2+) levels. May also regulate potassium ion transport by phosphorylation of potassium channel subunits. Phosphorylates SRC; this increases SRC kinase activity. Phosphorylates ASAP1, NPHP1, KCNA2 and SHC1. Promotes phosphorylation of ASAP2, RHOU and PXN; this requires both SRC and PTK2/PYK2 (By similarity).  
[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) uses an alternate in-frame splice site in the 3' coding region compared to variant 1. This results in a shorter protein (isoform 2) compared to isoform 1.