

Product datasheet for **MC223364**

Pik3cd (NM_001164051) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Pik3cd (NM_001164051) Mouse Untagged Clone
Tag: Tag Free
Symbol: Pik3cd
Synonyms: 2410099E07Rik; 2610208K16Rik; AW545373; p110delta
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC223364 representing NM_001164051
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGCCCCCTGGGGTGGACTGCCCATGGAGTTCTGGACCAAGAGGAGAGCCAGAGCGTGGTTGTTGACT
 TCTTGCTGCCACAGGGGTCTACTTGAACCTCCCGTGTCCCGCAATGCCAACCTCAGCACCATCAAGCA
 GGTGCTGTGGCACCCTGCACAGTATGAGCCACTTCCACATGCTCAGTGACCCCGAGGCCTATGTGTT
 ACCTGTGTGAACCAGACGGCGGAGCAGCAGGAGTTGGAGGATGAGCAGCGGAGGCTGTGCGACATCCAGC
 CCTTCTGCCCGTGTGCGCCTCGTGGCCCGAGAGGGGACCGCGTGAAGAAGCTCATTAACTCCAGAT
 CAGCCTCCTCATTGGCAAAGGTCTCCATGAGTTTGATTCCCTGCGGGACCCGGAAGTAAACGACTCCGC
 ACTAAGATGCGCCAGTTTTGTGAAGAGGCTGCTGCTCACCGCCAGCAGCTGGGCTGGGTGGAATGGCTGC
 AGTACAGCTTCCCGTGCAGCTGGAGCCCTCAGCAAGGGGTTGGCGGGCCGGCTTATTGCGTGTGAGCA
 CCGAGCCCTGTGGTCAACGTGAAGTTCGAGGGCAGTGGAGAGCTTCCCTCCAGGTATCCACCAAG
 GACATGCCCTGGCACTGATGGCCTGTGCCCTCCGAAAAAGGCCACAGTGTCCGGCAGCCTCTGGTGG
 AGCAGCCTGAGGAATATGCCCTGCAGTGAACGGGAGGCACGAATACCTCTACGGCACTACCCGCTCTG
 CCACTTTTCAGTACATCTGCAGCTGCCTACACAGCGGGCTGACCCCTCATCTGACCATGGTCCACTCCTC
 TCCATCCTTGCTATGCGGGATGAGCAGAGCAATCCTGCCCCCAAGTACAGAAACCACGTGCCAAACCTC
 CCCCAGTCCCTGCCAAGAAGCCCTCCTCTGTGTCCCTGTGGTCCCTGGAACAGCCATTCTCCATTGAGCT
 GATCGAGGGCCGAAAAAGTGAATGCTGACGAGCGGATGAAGCTGGTTGTTGAGCCGGGCTCTTCCATGGC
 AATGAGATGCTGTGCAAGACTGTGTCAAGCTCGGAGGTGAATGTATGCTCAGAGCCCGTGTGGAAGCAGC
 GACTGGAGTTCGATATCAGCGTCTGTGACCTCCCGCGCATGGCTCGACTCTGTTTTGCTCTATGCCGT
 CGTGGAGAAGGCTAAGAAGGCACGCTCCACAAAGAAGAAGTCTAAGAAGGCGGACTGCCCATCGCTTGG
 GCCAACCTCATGTATTGACTACAAAGATCAGCTCAAGACGGGGAGCGCTGCCTCTACATGTGGCCCT
 CTGTCCCAGATGAGAAGGGAGAGCTGCTGAATCCTGCGGGTACAGTGCCGGGGAACCCCAACACGGAGAG
 TGCCGCTGCCCTGGTCTACCTGCCTGAGGTGGCCCCCACCCTGTGTACTTCCCGCTCTGGAGAAG
 ATCCTGGAGCTGGGGCGTCACGGGGAGCGTGGGCGCATACGGAGGAGGAGCTGCAGCTGCGGGAGATCC



TGGAACGGCGGGGATCCGGGAACTGTACGAACATGAGAAGGACCTGGTGTGGAAGATGCGCCACGAAGT
 CCAGGAGCATTTCCAGAGGCGCTGGCCCGCTGCTGCTGGTACCAAGTGAATAAACACGAGGATGTG
 GCCCAGCTGTCCAGATGCTCTATTTGCTGTGCTCCTGGCCGAGCTGCCTGTGCTGAGCGCCCTGGAAC
 TTCTGGACTTTAGCTTTCCGACTGCTACGTGGGCTCCTTCGCCATCAAGTCCCTTCGGAAGCTGACGGA
 CGATGAGCTCTCCAGTACCTTTCGAGCTGGTGAAGTGTCAAATATGAGTCTACCTGGACTGCGAG
 CTGACCAAATCTTGTGGGCGGAGCCCTGGCTAACCGCAAGATCGGACACTTCTGTTCTGGCACCTCC
 GCTCTGAGATGCACGTACCATCAGTGGCTCTGCGGTTTGGTCTCATCATGGAAGCCTACTGCAGAGGCAG
 CACCCACCACATGAAGGTGCTGATGAAGCAGGGGGAAGCACTGAGCAAGCTTAAGGCACTGAATGACTTT
 GTGAAGGTGAGTTCCAGAAAGACCACCAAGCCCAAAACCAAGGAGATGATGCATATGTGCATGCGCCAGG
 AGACCTACATGGAGGCCCTGTCCACCTGCAGTCTCCACTCGACCCAGCACCCCTGCTGGAGGAAGTCTG
 TGTGGAGCAGTGACCTTCATGGACTCCAAAATGAAGCCCTGTGGATCATGTACAGCAGCGAGGAGGCG
 GGCAGTGTGGCAACGTGGGCATCATCTTAAGAACGGGGATGACCTCCGCCAGGACATGCTGACTCTGC
 AGATGATCCAGCTCATGGACGTCTGTGGAAGCAGGAGGGCTGGACCTGAGGATGACGCCCTACGGCTG
 CCTCCCCACGGGACCCACAGGTCTCATCGAGGTGGTCTCCACTCGGACACCATCGCCAACATCCAG
 CTGAACAAAAGCAACATGGCGGCCACAGCTGCCTTCAACAAGGACGCCCTGCTCAACTGGCTCAAGTCCA
 AGAACCTGGGGAGGCCCTGGATCGGGCATTGAGGAATTCACCTCTCCTGTGCTGGCTACTGTGTGGC
 CACATATGTTCTGGGCATCGGTGACCGGCACAGCGACAACATCATGATCAGAGAGAGTGGGCAGCTCTTC
 CACATTGATTTTGGCCACTTTCTGGGAACTTCAAGACCAAGTTTGAATCAACCGAGAGCGCGTCCCTT
 TCATTCTACCTACGACTTTGTCCACGTGATCCAGCAGGGGAAGACTAACCAACAGTGAGAAGTTTGAAG
 GTTCCGCGGCTACTGTGAACGAGCCTATACCATCTGCGGCGCCACGGGCTGCTTTTCTCCATCTCTTC
 GCCCTGATGCGGGCCGAGGTCTGCCTGAGCTTAGCTGCTCCAAGATATCCAGTATCTCAAGGACTCTC
 TGGCACTGGGGAAGACGGAGGAAGAGGGCCTAAAGCACTTCCGGGTGAAGTTCAACGAAGCTCTCCGAGA
 AAGCTGGAAAACCAAGTCAACTGGCTGGCGACAATGTGTCCAAGGATAACCGACAGTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:

Sgfl-Mlul

ACCN:

NM_001164051

Insert Size:

3141 bp

OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq:

NM_001164051.1, NP_001157523.1

RefSeq Size:

4942 bp

RefSeq ORF:

3141 bp

Locus ID: 18707

Cytogenetics: 4 E2

Gene Summary: Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns(4,5)P₂ (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP₃). PIP₃ plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Mediates immune responses. Plays a role in B-cell development, proliferation, migration, and function. Required for B-cell receptor (BCR) signaling. Mediates B-cell proliferation response to anti-IgM, anti-CD40 and IL4 stimulation. Promotes cytokine production in response to TLR4 and TLR9. Required for antibody class switch mediated by TLR9. Involved in the antigen presentation function of B-cells. Involved in B-cell chemotaxis in response to CXCL13 and sphingosine 1-phosphate (S1P). Required for proliferation, signaling and cytokine production of naive, effector and memory T-cells. Required for T-cell receptor (TCR) signaling. Mediates TCR signaling events at the immune synapse. Activation by TCR leads to antigen-dependent memory T-cell migration and retention to antigenic tissues. Together with PIK3CG participates in T-cell development. Contributes to T-helper cell expansion and differentiation. Required for T-cell migration mediated by homing receptors SELL/CD62L, CCR7 and S1PR1 and antigen dependent recruitment of T-cells. Together with PIK3CG is involved in natural killer (NK) cell development and migration towards the sites of inflammation. Participates in NK cell receptor activation. Have a role in NK cell maturation and cytokine production. Together with PIK3CG is involved in neutrophil chemotaxis and extravasation. Together with PIK3CG participates in neutrophil respiratory burst. Have important roles in mast-cell development and mast cell mediated allergic response. Involved in stem cell factor (SCF)-mediated proliferation, adhesion and migration. Required for allergen-IgE-induced degranulation and cytokine release. The lipid kinase activity is required for its biological function.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (6) represents use of an alternate promoter and 5' UTR, and uses two alternate in-frame splice sites in the central coding region, compared to variant 4. The resulting isoform (e) lacks a 1-aa internal segment and includes an additional 3-aa internal segments, compared to isoform c. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.