

Product datasheet for **MC223607**

Upf1 (NM_030680) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Upf1 (NM_030680) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Upf1
Synonyms:	B430202H16Rik; NORF1; PNORF-1; Rent1; Upflp
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
Fully Sequenced ORF:	>MC223607 representing NM_030680 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGAGTGTGGAGGCGTACGGCCCCAGCTCGAAACACTCACCTTCTTGGACTGAGGAGCCGAGCTGC
TCGGCGCCGACACCCAGGGCTCCGAGTTCGAATTCACCGACTTACCCTTCCCAGCCAGACGCAGACGCC
CCCCGGCGGCCCGGGCGCGGGAGGCCCGGGCGGAGCGGGCGCAGGCGCGCGGCCAGCTCGAC
GCACAAGTTGGACCAGAGGGCATCTTGCAAAATGGGGCTGTGGATGACAGTGTGGCAAGACCAGCCAGC
TGCTAGCTGAGCTGAACCTCGAGGAAGATGAAGAGGACACATACTACACTAAGGACCTCCCAGTCCACGC
CTGCAGTTACTGTGGAATCCATGATCCTGCCTGCGTGGTTTACTGTAATACCAGCAAGAAGTGGTTCTGC
AATGGCCGAGGAAATACTTCTGGCAGCCACATTGTGAATCACCTCGTGAGGGCAAAATGCAAGGAAGTGA
CGCTGCACAAGGACGGCCCTCTGGGCGAGACCGTGTGGAGTGTACAACCTGTGGCTGCCCAACGTCTT
CCTGCTGGGCTTCATCCCTGCGAAGGCCGACTCTGTGGTGGTGTGTTGTGCAGGCAGCCCTGTGCCAGC
CAGAGCAGCCTGAAGGACATCAACTGGGACAGCTCACAGTGGCAGCCCTAATCCAGGACCGTGTCTTTC
TGTCATGGCTGGTCAAGATTCCCTCTGAGCAGGAGCAGCTGCGAGCACGGCAGATCACGGCACAGCAGAT
CAACAAGCTGGAAGAGCTCTGGAAGGAAAATCCTTCAGCCACTCTGGAGGACCTGGAGAAGCCAGGCGTA
GACGAGGAGCCACAGCACGTGCTCCTGCGTTACGAGGATGCTTACCAGTACCAGAACATCTTCGGGCCAC
TGGTCAAGCTGGAGGCTGACTATGACAAGAAGTTGAAGGAGTACAGACTCAAGATAACATCACGGTCAG
GTGGGACCTGGGCCTTAACAAGAAGAGAATCGCCTTCTTCACTTTGCCAAGACTGACTCTGACATGCGG
CTCATGCAGGGTATGAGATCTGTCTGCGGTACAAAGGGGATCTGGCGCCCTGTGGAAGGGGATTGGCC
ACGTATCAAGGTTCTGATAATTATGGTATGAGATTGCTATTGAGCTCCGAGCAGCGTGGTGGCCCC
TGTGGAAGTGACCCACAACCTCCAAGTGGATTTGTGTGGAAGTCAACCTCTTTTGATAGGATGCAGAGT
GCACTGAAGACCTTCGCTGTGGACGAGACCTCTGTGTCAGGGTATATTTACCACAAGCTGCTGGGCCAGC
AGGTGGAGGATGTGGTCAAGTGCCAGCTGCCAAAGCGCTTCACAGCTCAGGGGCTCCCTGACCTCAA
CCACTCTCAGGTGATGCTGTGAAGACCGTGTGTCAGAGACCACTCAGCCTCATCCAGGGCCCTCCAGGC
ACAGGCAAGACTGTGACATCAGCCACTATTGTCTACCACCTTGTCTCGGCAGGGCAATGGGCCTGTACTGG



[View online »](#)

```

TTTGTGCTCCAAGTAACATCGCTGTGGACCAGCTCACAGAGAAGATCCACCAGACAGGACTGAAGGTCGT
ACGCCTCTGTGCCAAGAGCCGTGAGGCCATTGACTCCCCAGTGTCTTCCCTGGCTTTGCACAACCAGATC
AGGAACATGGACAGCATGCTGAGCTGCAGAAGCTGCAGCAGCTAAAGGATGAGACAGGCGAGCTGCAT
CTGCAGATGAGAAGCGGTACCGGGCGCTTAAGCGCACAGCTGAGAGAGAAGTCTCATGAATGCAGATGT
CATATGCTGCACATGTGTGGTGTGGTACCCGAGGCTGGCCAAGATGCAGTCCGTTCCATCCTCATT
GACGAGAGCACCCAGGCCACTGAGCCTGAGTGCATGGTGCCTGTAGTCTTGGGGCCAAGCAGCTAATCC
TCGTCCGTGACCACTGCCAGCTGGGCCAGTGGTGTGATGTGCAAGAAGGCAGCCAAGGCCGACTGTACA
ATCGCTCTTCGAGCGCTTGGTGGTGTGGGCATCCGGCCATCCGCTGCAGGTGCAATACCGCATGCAC
CCTGCACTCAGCGCCTTCCGTCCAACATCTTCTACGAGGGCTCATTGCAGAATGGCGTCACTGCAGCGG
ATCGTGTCAAAAAAGGCTTTGACTTCCAGTGGCCACAACCTGACAAGCCTATGTTCTTCTACGTGACGCA
GGGCCAGGAGGATTGCCAGCTCTGGCACATCTACCTCAACAGGACGGAGGCAGCCAATGTGGAGAAG
ATAACTACGAAGCTGTTGAAGGCAGGTGCAAGCCTGACCAGATCGGCATCATCACCCCCACGAGGGCC
AGCGCTTACTTGGTGCAGTACATGCAGTTCAGCGGCTCCCTGCACACAAAGCTCTACCAGGAAGTGA
AATTGCCAGTGTGACGCCTTCCAGGGCCGGGAGAAGGACTTCATATTCTGTCTCGCTGCGCGCCAAT
GAACATCAGGGCATTGGGTTCTAAACGACCCCGGCGTCTGAATGTGGCTCTCACCAGAGCAAGATATG
GGTGATCATTGTGGTAACCCAAAGGCCCTGTGGAAGCAGCCCTGTGGAATCAGCTGAGCTACTA
CAAGGAACAGAAGGCGCTAGTGAAGGGCCGCTCAACAACCTACGTGAGAGCCTCATGCAGTTCAGCAAG
CCTCGAAACTTGTCAACACTGTCAACCCGGGTGCCCGCTTATGACTACTGCCATGTACGATGCCCGTG
AGGCCATCATCCCGGTCTGTCTATGACCGCAGCAGCCAGGGCCGGCCCTCGAACATGTACTTCCAGAC
CCATGACCAGATTAGTATGATCAGCGCAGGCCACCGCCACGTGGCTGCCATGAACATCCCTATTCCCTTC
AACTTGGTCATGCCTCCCATGCCCCACCTGGCTACTTCGGACAGGCCAACGGCCGGCAGCTGGTGGG
GCACCCCAAAAACCAAGACTGGCCGTGGGGCCGCCAGAAGAACCCTTTGGGCTTCTGGGCCAGCCA
GACCACCTTCCCAACAGCCAGGCCAGGACAGTGGCCTCCAGCCCTTTTACAGGGTGCCCTCACA
CAGGGTTACGTGTCCATGAGCCAGCCCTCTCAGATGAGCCAGCCCTGGCCCTCTCCAGCCAGAAGTGTCCC
AGGACAGTACCTCGGTGATGAGTTTAAATCACAGATTGACGTGGCACTCTCACAAGACTCCACATACCA
GGGAGAGCGGGCATACCAGCACGGCGGGTACCAGGGCTGTCCAGTACTAG

```

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM_030680
- Insert Size:** 3342 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
- RefSeq:** [NM_030680.2](#), [NP_109605.2](#)

RefSeq Size: 4598 bp

RefSeq ORF: 3342 bp

Locus ID: 19704

UniProt ID: [Q9EPU0](#)

Cytogenetics: 8 34.15 cM

Gene Summary: RNA-dependent helicase and ATPase required for nonsense-mediated decay (NMD) of mRNAs containing premature stop codons. Is recruited to mRNAs upon translation termination and undergoes a cycle of phosphorylation and dephosphorylation; its phosphorylation appears to be a key step in NMD. Recruited by release factors to stalled ribosomes together with the SMG1C protein kinase complex to form the transient SURF (SMG1-UPF1-eRF1-eRF3) complex. In EJC-dependent NMD, the SURF complex associates with the exon junction complex (EJC) (located 50-55 or more nucleotides downstream from the termination codon) through UPF2 and allows the formation of an UPF1-UPF2-UPF3 surveillance complex which is believed to activate NMD. Phosphorylated UPF1 is recognized by EST1B/SMG5, SMG6 and SMG7 which are thought to provide a link to the mRNA degradation machinery involving exonucleolytic and endonucleolytic pathways, and to serve as adapters to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation. UPF1 can also activate NMD without UPF2 or UPF3, and in the absence of the NMD-enhancing downstream EJC indicative for alternative NMD pathways. Plays a role in replication-dependent histone mRNA degradation at the end of phase S; the function is independent of UPF2. For the recognition of premature termination codons (PTC) and initiation of NMD a competitive interaction between UPF1 and PABPC1 with the ribosome-bound release factors is proposed. The ATPase activity of UPF1 is required for disassembly of mRNPs undergoing NMD (By similarity). Essential for embryonic viability.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) uses an alternate in-frame splice junction compared to variant 1. The resulting isoform (b) is shorter compared to isoform a.