

## Product datasheet for MC223689

### Dab2ip (NM\_001114125) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Dab2ip (NM\_001114125) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Dab2ip  
**Synonyms:** 2310011D08Rik; AI480459; Aip1; mKIAA1743  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**Fully Sequenced ORF:** >MC223689 representing NM\_001114125  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCCGCGATCGCC

ATGGAGCCCGACTCCCTCCTGGACCCAGGGGACTCCTACGAGTCGCCCAAGAAAGGCCGGGCTCCCGC  
 GGAGCCTTCCCGGCAGCATGTCTAGAGAAAAACCCAGCATGGAGCCCTCGGCTTCAACCCGTTCCGGGT  
 CACGGGCTTCTCAGCCGCCCTCAAGGGCTCCATCAAGCGCACCAAGGCCAGCCAAACTGGACCGC  
 AACCCACAGCTTCCGCCACATCCTGCCGGGTTCGGAGCGCAGCCGCCCGCCGCGGACAATGAGAGGT  
 CCCATCTGATGCCAAGGCTGAAGGAGTCTCGGTACACGAGTCCCTGCTCAGCCCCAGCAGCGCAGTGG  
 GGCCCTGGACCTCAGCATGGAGGAGGAGGTGATTATCAAGCCCGTTCACAGCAGCATCCTGGGTCAGGAC  
 TACTGCTTCGAGGTGACAACATCATCAGGAAGCAAGTGTTCCTCGCCGGTTCAGCCGCTGAGCGCGATA  
 AGTGGATGGAGAACCTGAGGCGAGCAGTGCACCCCAACAAGGACAACAGCCGGCGTGTGGAGCATCCT  
 GAAGCTGTGGGTGATTGAGGCCAAGGATCTGCCGGCAAGAAGAAGTATCTATGTGAAGTGTGCCTGGAC  
 GATGTGCTGTATGCCGTACCACAAGCAAGCTCAAGACGGACAATGTCTTCTGGGAGAGCACTTTGAGT  
 TCCATAACCTGCCGCTCTACGCACAGTCACTGTGCACCTGTATCGGGAGACTGACAAGAAAAAGAAAA  
 GGAACGCAACAGTACCTGGGCCTGGTGAGCCTGCCTGCCGCTCTGTGGCTGGGCGGCAGTTTGTGGAG  
 AAGTGGTACCCAGTGGTACACCCCAACCCCAAGGGTGGCAAGGCCCTGGGCCCATGATCCGAATCAAGG  
 CACGCTACCAGACCGTCAGCATCTTGCCATGAGAGATGTACAAGGAGTTTGGCGAGCACATCACTAACCA  
 CTACCTGGGGCTGTGCGCAGCCCTGGAACCCATCCTCAGTGCCAAGACCAAGGAGGAGATGGCGTCGGCT  
 CTGGTGACATCCTGCAGAGCACGGAAAGGTGAAGGACTTTCTAACAGACCTGATGATGCAGAGGTGG  
 ACCGCTGTGGGACAATGAGCACCTCATCTTCCGGGAGAACACTGGCCACCAAGGCCATCGAGGAATA  
 CCTCAAATTTGTGGCCAGAAGTACCTGCAGGACGCACTAGGTGAGTTCATCAAAGCTCTGTATGAGTCA  
 GATGAAAATTGTGAAGTGGACCAAGCAAGTGCTCATCCGCTGACCTCCCTGAGCACCAGGGCAACCTCA  
 AGATGTGCTGTGAGCTGGCCTTCTGCAAGATCATCAACTCTACTGCGTCTTCCACGGGAGCTTAAGGA  
 GGTGTTCCCTCATGGCGGCAGGAGTGTAGCAGCCGAGGCCGAGATATCAGTGAACGGCTCATCAGC  
 GCCTCCCTCTTCTCGCTTCTGTGCCCTGCCATCATGTACCCTCGCTCTTCAACTGCTTCAGGAGT



[View online »](#)

```

ATCCTGACGACCGCACGGCTCGCACCTCACGCTCATTGCCAAAGTCACCCAGAACCTGGCCAACTTTGC
CAAGTTTGGCAGCAAGGAAGAATACATGTCTTTCATGAACCAGTTCCTGGAGCACGAGTGGACCAACATG
CAGCGCTTCTGTTGGAGATCTCCAACCCCGAGACCTTTCCAACACAGCAGGCTTCGAGGGCTACATAG
ACCTGGGCGGGGAGCTCTAGCCTGCACTCCCTGCTCTGGGAAGCTGTCAGCCAGCTTGATCAGAGCGT
TGTGTCGAAGCTGGGACCTCTGCCTCGTATCCTGAGGGATGTCCACACAGCACTGAGCACTCCTGGCAGT
GGGCAGCTCCCTGGCACCATGACCTGCCTCCACCCCGGGCTCCGGCAGCAGCAGCGTCTCTGCTGGGC
TTCAGAAGATGGTGATTGAAAATGACCTCTCTGGTCTGATAGATTTACCCCGGTTACCGTCTCCAACCCC
CGAAAACAAGGACTTGTTTTTGTCAAAAGTCTCCGGGGTCCAGCCTTCACTGCCCGCAGCTCAAGC
TACTCAGAAGCCAATGAACCTGACCTGCAGATGGCCAATGGCAGCAAGAGCCTGTCCATGGTGGACCTCC
AGGACGCCCGCACGCTGGATGGGAGGCAGGTTCCCCAGTGGGCCAGACGCCCTACCTGCTGACGGGCA
GGTGCCTGCGACTCAGCTGCTGGTGGGAGCCAGGCAGCCAGTGGCAGCTGGCAGGATTGGCC
ACAGTGCAGCGGGCAGTGCCAACACCAACCACACCAGGCACCTCCGAGGGTGCACCAGGACGGCCCCAGT
TGTTGGCCCCACTTTCCTCCAGAATCCTGTGTACCAGATGGCGGCCGGCCTGCCACTGTCACCCCGTGG
CCTTGGTGACTCAGGCTCTGAAGGCCACAGCTCCCTGAGCTCTCACAGCAACAGTGAAGAGCTGGCAGCC
GCTGCCAAACTAGGAAGTTTCAGCACTGCTGCAGAGGAGCTGGCAAGGGCGCCTGGAGAAGTGGCACGGA
GGCAGATGCTCACTGACTGAGAAGGGTGGCAGCCACAGTGCAGAGGCAAAATAGTGCCGGTCCCCAGCG
GAGGATTGACCAGCCGCCACCGCCACCACCACCACCGCCTCCTGCTCCCCGGGGCAGGACACCTCCTACC
CTGCTGAGCACCTACAGTACCCACGACCTCAAGTGGAAACCCTGGCATCAGCATCCCCGACTGGGCTG
GCCCTGGCACCCGGCTGCGGCAACAGTCTCCTCCTCAAAGGGAGACAGCCAGAGCTGAAGCCCCGAGC
CATGCACAAGCAGGGCCCTTACCCGTCAGTCCAATGCCCTGGACCGCACGGCCGCTTGGCTCTTGACC
ATGAACCGCAGTTGTTAGAAGACGAGGGTCTGGGCCAGATCCCCCCACAGGGATAGGCTAAGGAGTA
AGGAGGAACTCAGCCAAGCAGAAAAGGATCTGGCAGTGTACAAGACAAGCTACGGATCTCCACCAAGAA
GCTGGAGGAGTATGAGACCCTATTCAAGTGCAGGAGGAGACGACGAGCAAGCTGGTGGTATCAG
GCTCGGCTGGAAGAAGGTGAGGAGCGGCTGCGGGCAGCAGGAAGACAAGGATATCCAGATGAAAAGGCA
TCATCAGCAGTTGATGTGAGTGAAGAAGAAGTGAAGAAGGATCATGCAGAGATGCAAGCAGCTGTAGA
TTCCAACAGAAGATCATCGATGCCAGGAAAAGCGCATTGCCTCGTGGATGCTGCCAATGCCCGCCTC
ATGAGTGCCCTCACACAGCTGAAAGAGAGTATGCATTAG
    
```

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

- Restriction Sites:** SgfI-MluI
- ACCN:** NM\_001114125
- Insert Size:** 3399 bp
- OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
- Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
  2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
  3. Close the tube and incubate for 10 minutes at room temperature.
  4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
  5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: NM\_001114125.1, NP\_001107597.1  
RefSeq Size: 5388 bp  
RefSeq ORF: 3399 bp  
Locus ID: 69601  
UniProt ID: Q3UHC7  
Cytogenetics: 2 B

**Gene Summary:**

Functions as a scaffold protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways. Involved in several processes such as innate immune response, inflammation and cell growth inhibition, apoptosis, cell survival, angiogenesis, cell migration and maturation. Plays also a role in cell cycle checkpoint control; reduces G1 phase cyclin levels resulting in G0/G1 cell cycle arrest. Mediates signal transduction by receptor-mediated inflammatory signals, such as the tumor necrosis factor (TNF), interferon (IFN) or lipopolysaccharide (LPS). Modulates the balance between phosphatidylinositol 3-kinase (PI3K)-AKT-mediated cell survival and apoptosis stimulated kinase (MAP3K5)-JNK signaling pathways; sequesters both AKT1 and MAP3K5 and counterbalances the activity of each kinase by modulating their phosphorylation status in response to proinflammatory stimuli. Acts as a regulator of the endoplasmic reticulum (ER) unfolded protein response (UPR) pathway; specifically involved in transduction of the ER stress-response to the JNK cascade through ERN1. Mediates TNF-alpha-induced apoptosis activation by facilitating dissociation of inhibitor 14-3-3 from MAP3K5; recruits the PP2A phosphatase complex which dephosphorylates MAP3K5 on 'Ser-966', leading to the dissociation of 13-3-3 proteins and activation of the MAP3K5-JNK signaling pathway in endothelial cells. Mediates also TNF/TRAF2-induced MAP3K5-JNK activation, while it inhibits CHUK-NF-kappa-B signaling. Acts a negative regulator in the IFN-gamma-mediated JAK-STAT signaling cascade by inhibiting smooth muscle cell (VSMCs) proliferation and intimal expansion, and thus, prevents graft arteriosclerosis (GA). Acts as a GTPase-activating protein (GAP) for the ADP ribosylation factor 6 (ARF6) and Ras. Promotes hydrolysis of the ARF6-bound GTP and thus, negatively regulates phosphatidylinositol 4,5-bisphosphate (PIP2)-dependent TLR4-TIRAP-MyD88 and NF-kappa-B signaling pathways in endothelial cells in response to lipopolysaccharides (LPS). Binds specifically to phosphatidylinositol 4-phosphate (PtdIns4P) and phosphatidylinositol 3-phosphate (PtdIns3P). In response to vascular endothelial growth factor (VEGFA), acts as a negative regulator of the VEGFR2-PI3K-mediated angiogenic signaling pathway by inhibiting endothelial cell migration and tube formation. In the developing brain, promotes both the transition from the multipolar to the bipolar stage and the radial migration of cortical neurons from the ventricular zone toward the superficial layer of the neocortex in a glial-dependent locomotion process. Probable downstream effector of the Reelin signaling pathway; promotes Purkinje cell (PC) dendrites development and formation of cerebellar synapses. Functions also as a tumor suppressor protein in prostate cancer progression; prevents cell proliferation and epithelial-to-mesenchymal transition (EMT) through activation of the glycogen synthase kinase-3 beta (GSK3B)-induced beta-catenin and inhibition of PI3K-AKT and Ras-MAPK survival downstream signaling cascades, respectively.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) contains an alternate 5' terminal exon, and it thus differs in the 5' UTR and initiates translation at an alternate start codon, and it also lacks a 3' segment that results in an alternate 3' coding region, compared to variant 2. The encoded isoform (1) has distinct N- and C-termini and is shorter than isoform 2. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.