

Product datasheet for **MC224837**

Siglec1 (NM_011426) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Siglec1 (NM_011426) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Siglec1
Synonyms:	Cd169; Siglec-1; Sn
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_011426
Insert Size:	5106 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_011426.3 , NP_035556.3
RefSeq Size:	6427 bp
RefSeq ORF:	5106 bp



[View online »](#)

Locus ID: 20612

UniProt ID: [Q62230](#)

Cytogenetics: 2 63.26 cM

Gene Summary: Acts as an endocytic receptor mediating clathrin dependent endocytosis. Macrophage-restricted adhesion molecule that mediates sialic-acid dependent binding to lymphocytes, including granulocytes, monocytes, natural killer cells, B-cells and CD8 T-cells (By similarity). Preferentially binds to alpha-2,3-linked sialic acid. Binds to SPN/CD43 on T-cells. May play a role in hematopoiesis. May act as a counter-receptor for CLEC10A in lymph node.
[UniProtKB/Swiss-Prot Function]