

## **Product datasheet for MC225086**

## Notch4 (NM\_010929) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Notch4 (NM\_010929) Mouse Untagged Clone

Tag: Tag Free Symbol: Notch4

Synonyms: Int-3; Int3; N4

Mammalian Cell

Selection:

**Vector:** pCMV6-Entry (PS100001)

Neomycin

E. coli Selection: Kanamycin (25 ug/mL)

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_010929

**Insert Size:** 5895 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20  $^{\circ}$ C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 010929.2, NP 035059.2

RefSeq Size: 6604 bp RefSeq ORF: 5895 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Notch4 (NM\_010929) Mouse Untagged Clone - MC225086

**Locus ID:** 18132

UniProt ID: P31695

Cytogenetics: 17 18.15 cM

**Gene Summary:** Functions as a receptor for membrane-bound ligands Jagged1, Jagged2 and Delta1 to

regulate cell-fate determination. Upon ligand activation through the released notch intracellular domain (NICD) it forms a transcriptional activator complex with RBPJ/RBPSUH

and activates genes of the enhancer of split locus. Affects the implementation of

differentiation, proliferation and apoptotic programs (By similarity). May regulate branching

morphogenesis in the developing vascular system.[UniProtKB/Swiss-Prot Function]