

Product datasheet for MC225484

Spink2 (NM_001289768) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Spink2 (NM_001289768) Mouse Untagged Clone
Tag: Tag Free
Symbol: Spink2
Synonyms: 1700007F22Rik; AV038945; HUSI-II
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Fully Sequenced ORF: >MC225484 representing NM_001289768

Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGCTGAGACTGGTGCTGTTGCTCCTGGTCACAGACTTTGCAGCCTCTCATGAGACTCTCGACTCTTCCG
 ATTCTCAAATCATGAAGAGGTCACAGTTCCGAACAACGTGGTCATTTGACTTCCCAGCATGCCCTAGG
 AACCTCAACCCTGTGTGCGGAACGGATATGA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001289768
Insert Size: 171 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.



[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001289768.1, NP_001276697.1</u>
RefSeq Size:	627 bp
RefSeq ORF:	171 bp
Locus ID:	69982
UniProt ID:	<u>Q8BMY7</u>
Cytogenetics:	5 C3.3
Gene Summary:	<p>As a strong inhibitor of acrosin, it is required for normal spermiogenesis. It probably hinders premature activation of proacrosin and other proteases, thus preventing the cascade of events leading to spermiogenesis defects (PubMed:21705336, PubMed:28554943). May be involved in the regulation of serine protease-dependent germ cell apoptosis (PubMed:21705336). It also inhibits trypsin (PubMed:21705336).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (4) differs in the 5' UTR and 5' coding region and uses an alternate splice site in the central coding region, resulting in a frameshift and an early stop codon, compared to variant 1. It encodes isoform 4, which is shorter and has distinct N- and C-termini, compared to isoform 1.</p>