

Product datasheet for MC225683

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Sh3kbp1 (NM_001290661) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Sh3kbp1 (NM 001290661) Mouse Untagged Clone

Tag: Tag Free Symbol: Sh3kbp1

Synonyms: 1200007H22Rik; 1700125L08Rik; 5830464D22Rik; Al447724; Cin85; IN85; Ruk; Seta

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC225683 representing NM_001290661

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001290661

Insert Size: 333 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.



Sh3kbp1 (NM_001290661) Mouse Untagged Clone - MC225683

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001290661.1</u>, <u>NP 001277590.1</u>

RefSeq Size: 3850 bp
RefSeq ORF: 333 bp
Locus ID: 58194
UniProt ID: Q8R550
Cytogenetics: X F4

Gene Summary: Adapter protein involved in regulating diverse signal transduction pathways. Involved in the

regulation of endocytosis and lysosomal degradation of ligand-induced receptor tyrosine kinases, including EGFR and MET/hepatocyte growth factor receptor, through an association with CBL and endophilins. The association with CBL, and thus the receptor internalization, may be inhibited by an interaction with PDCD6IP and/or SPRY2. Involved in regulation of ligand-dependent endocytosis of the IgE receptor. Attenuates phosphatidylinositol 3-kinase activity by interaction with its regulatory subunit. May be involved in regulation of cell adhesion; promotes the interaction between TTK2B and PDCD6IP. May be involved in the

regulation of cellular stress response via the MAPK pathways through its interaction with MAP3K4. Is involved in modulation of tumor necrosis factor mediated apoptosis. Plays a role

in the regulation of cell morphology and cytoskeletal organization. Required in the control of

cell shape and migration (By similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4, also known as Rukh2) lacks multiple coding exons and contains an alternate 5' exon, compared to variant 1. It initiates translation at a downstream in-frame start codon. The encoded isoform (4) has a shorter N-terminus than isoform 1.

Variants 4 and 5 encode the same isoform (4).