

## Product datasheet for MC225684

## Sh3kbp1 (NM\_001290664) Mouse Untagged Clone

## **Product data:**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Sh3kbp1 (NM_001290664) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Sh3kbp1
Synonyms:	1200007H22Rik; 1700125L08Rik; 5830464D22Rik; Al447724; Cin85; IN85; Ruk; Seta
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC225684 representing NM_001290664 <mark>Red=</mark> Cloning site Blue=ORF Orange=Stop codon
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGCAGCTGCCAGCAGTGGGCCAGCTTCTCTCTCTCTCAGTGGCATCCTCACCCATGTCATCCTCTTTGG GAACAGCTGGACAGAGAGCCAGTTCTCCATCTCTGTTCAGCACAGAAGGAAAGCCAAAGATGGAGCCAGC AGTGAGCAGCCAGGCTGCTATCGAGGAGCTTAAGATGCAAGTCCGTGAGGCCGAGGACCATCATTGAGACC ATGAAGGACCAGCAGAAACGTGAGATTAAGCAGTTACTGTCAGAATTGGATGAAGAGAAAAAGATCCGGC TCCGGTTGCAGATGGAAGTGAACGACATAAAGAAAGCTCTTCAATCAA
	ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAGGTTTAA
Chromatograms:	https://cdn.origene.com/chromatograms/ja3286_b09.zip
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001290664
Insert Size:	333 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Sh3kbp1 (NM_001290664) Mouse Untagged Clone – MC225684	
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Met	<ul> <li>hod: 1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ul>
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM 001290664.1, NP 001277593.1</u>
RefSeq Size:	3721 bp
RefSeq ORF:	333 bp
Locus ID:	58194
UniProt ID:	<u>Q8R550</u>
Cytogenetics:	X F4
Gene Summary:	Adapter protein involved in regulating diverse signal transduction pathways. Involved in the regulation of endocytosis and lysosomal degradation of ligand-induced receptor tyrosine kinases, including EGFR and MET/hepatocyte growth factor receptor, through an association with CBL and endophilins. The association with CBL, and thus the receptor internalization, may be inhibited by an interaction with PDCD6IP and/or SPRY2. Involved in regulation of ligand-dependent endocytosis of the IgE receptor. Attenuates phosphatidylinositol 3-kinase activity by interaction with its regulatory subunit. May be involved in regulation of cell adhesion; promotes the interaction between TTK2B and PDCD6IP. May be involved in the regulation of cellular stress response via the MAPK pathways through its interaction with MAP3K4. Is involved in modulation of tumor necrosis factor mediated apoptosis. Plays a role in the regulation of cell morphology and cytoskeletal organization. Required in the control of cell shape and migration (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (5, also known as Rukh4) lacks multiple coding exons and contains an alternate 5' exon, compared to variant 1. It initiates translation at a downstream in-frame start codon. The encoded isoform (4).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US