

Product datasheet for **MC225862**

II11 (NM_001290423) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	II11 (NM_001290423) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	II11
Synonyms:	IL-11
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001290423
Insert Size:	423 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001290423.1</u> , <u>NP_001277352.1</u>



[View online »](#)

RefSeq Size: 1845 bp

RefSeq ORF: 423 bp

Locus ID: 16156

Cytogenetics: 7 2.76 cM

Gene Summary: Cytokine that stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production (PubMed:8913282). Also promotes the proliferation of hepatocytes in response to liver damage (PubMed:22253262). Binding to its receptor formed by IL6ST and either IL11RA1 or IL11RA2 activates a signaling cascade that promotes cell proliferation, also in the context of various cancers (PubMed:10026196, PubMed:23948300). Signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3 (PubMed:23948300, PubMed:22253262).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) contains an alternate first exon compared to variant 1. The resulting isoform (2) is shorter at the N-terminus compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.