OriGene Technologies, Inc.
9620 Medical Center Drive, Ste 200
Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product datasheet for MC225892

Rbm4 (NM_001290124) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Rbm4 (NM_001290124) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Rbm4
Synonyms:	4921506I22Rik; lark; Lark1; Mlark; Rbm4a
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)

Fully Sequenced ORF:

```
>MC225892 representing NM_001290124
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCCGCGATCGCC

ATGGTGAAGCTGTTCATTGGAAATCTGCCCCGGGAGGCCACAGAGCAGGAGATCCGCTCACTCTTCGAGC
AGTACGGGAAGGTGCTGGAATGTGACATCATTAAGAATTATGGCTTTGTGCACATAGAGGACAAGACGGC
CGCTGAGGATGCCATACGCAACCTGCACCACTACAAGCTGCACGGAGTGAACATCAATGTGGAAGCCAGC
AAGAATAAGAGCAAAGCTTCAACCAAGTTACACGTGGGCAACATCAGCCCCACTTGTACCAACCAAGAGC
TTCGGGCCAAGTTTGAGGAGTACGGCCCAGTCATCGAATGTGACATCGTGAAAGATTATGCCTTTGTACA
CATGGAGCGGGCAGAGGATGCGGTGGAGGCCATCAGGGGCCTCGACAACACAGAGTTTCAAGGTGGGATG
TGTGTGGGCTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA
```

Restriction Sites:	SgfI-MluI
ACCN:	NM_001290124
Insert Size:	432 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.
©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001290124.1, NP_001277053.1
RefSeq Size:	1912 bp
RefSeq ORF:	432 bp
Locus ID:	19653
UniProt ID:	Q8C7Q4
Cytogenetics:	19 A
Gene Summary:	RNA-binding factor involved in multiple aspects of cellular processes like alternative splicing of pre-mRNA and translation regulation. Modulates alternative 5'-splice site and exon selection. Acts as a muscle cell differentiation-promoting factor. Activates exon skipping of the PTB pre-mRNA during muscle cell differentiation. Antagonizes the activity of the splicing factor PTBP1 to modulate muscle cell-specific exon selection of alpha tropomyosin. Binds to intronic pyrimidine-rich sequence of the TPM1 and MAPT pre-mRNAs. Required for the translational activation of PER1 mRNA in response to circadian clock. Binds directly to the 3' UTR of the PER1 mRNA. Exerts a suppressive activity on Cap-dependent translation via binding to CU-rich responsive elements within the 3' UTR of mRNAs, a process increased under stress conditions or during myocytes differentiation. Recruits EIF4A1 to stimulate IRES-dependent translation initiation in respons to cellular stress. Associates to internal ribosome entry segment (IRES) in target mRNA species under stress conditions. Plays a role for miRNA-guided RNA cleavage and translation suppression by promoting association of AGO2-containing miRNPs with their cognate target mRNAs. Associates with miRNAs during muscle cell differentiation. Binds preferentially to 5'-CGCGCG[GCA]-3' motif in vitro.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (4) differs in the 5' UTR, and lacks an alternate exon that results in a frameshift in the 3' coding region, compared to variant 1. The encoded isoform (b) has a distinct and shorter C-terminus, compared to isoform a. Both variants 4 and 5 encode isoform b.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use.
©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US