

Product datasheet for MC226126

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com

CN: techsupport@origene.cn

OriGene Technologies, Inc.

Srsf10 (NM_001284196) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Srsf10 (NM_001284196) Mouse Untagged Clone

Tag: Tag Free Symbol: Srsf10

Synonyms: Fusip1; Fusip1; Nssr; Nssr1; NssR2; Sfrs13a; Srrp40; Srsf13a; TASR; TASR1; TASR2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC226126 representing NM_001284196

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001284196

Insert Size: 549 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



Srsf10 (NM_001284196) Mouse Untagged Clone - MC226126

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001284196.1, NP 001271125.1</u>

4 D3

 RefSeq Size:
 3433 bp

 RefSeq ORF:
 549 bp

 Locus ID:
 14105

 UniProt ID:
 Q9R0U0

Cytogenetics:

Gene Summary: Splicing factor that in its dephosphorylated form acts as a general repressor of pre-mRNA

splicing. Seems to interfere with the U1 snRNP 5'-splice recognition of SNRNP70. Required for splicing repression in M-phase cells and after heat shock. Also acts as a splicing factor that specifically promotes exon skipping during alternative splicing. Interaction with YTHDC1, a RNA-binding protein that recognizes and binds N6-methyladenosine (m6A)-containing RNAs, prevents SRSF10 from binding to its mRNA-binding sites close to m6A-containing regions, leading to inhibit exon skipping during alternative splicing (By similarity). May be involved in regulation of alternative splicing in neurons (PubMed:10583508).[UniProtKB/Swiss-Prot

Function1

Transcript Variant: This variant (4) uses an alternate in-frame splice site in the 3' coding region and contains an alternate 3' terminal exon, compared to variant 2. It encodes isoform 4,

which is shorter and has a distinct C-terminus, compared to isoform 2.