

Product datasheet for MC226148

Cib1 (NM_001291275) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Cib1 (NM_001291275) Mouse Untagged Clone
Tag: Tag Free
Symbol: Cib1
Synonyms: Cibkip; Kip; Prkdcip
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC226148 representing NM_001291275
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGGAGGTTCCGGCAGTCGCCTGTCTAAGGAGCTGCTGGCCGAGTACCAGGACCTGACGTTCTGACCA
 AGCAGGAGATCCTCCTAGCCACAGACGCTTTTGTGAGCTGCTTCCCCAGAGCAGCGGACCGTGGAGGA
 GTCACTGCACACCCGAGTATCGTTTGAGCAGATCCTCAGCCTCCAGAGCTCAAGGAGCGAATCTGCATG
 GTCTTCTCCACATCACCTACCAGAGACAGCCTGAGCTTTGAGGACTTCTGGACCTCCTGAGTGTCTTCA
 GTGACACAGCAACCCAGACATCAAGTCACACTATGCCTTCCGCATCTTTGACTTTGATGACGATGGAAC
 CCTGGACAGAGAAGACCTGAGCCAGCTTGTGAATTGCCTCACGGGAGAGGGCGAGGACACTCGGCTCAGT
 GCTTCTGAGATGAAGCAGCTGATTGACAATATCCTGGAAGAGTCAGACATTGACAGGGATGGGACCATCA
 ATCTTTCCGAGTCCAGCATGTCATCTCGCGCTCACCAGACTTTGCCAGCTCCTTTAAGATTGTCCTGTG
 A

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001291275
Insert Size: 561 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



[View online »](#)

OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001291275.1</u> , <u>NP_001278204.1</u>
RefSeq Size:	1040 bp
RefSeq ORF:	561 bp
Locus ID:	23991
UniProt ID:	<u>Q9Z0F4</u>
Cytogenetics:	7 D2

Gene Summary:

Calcium-binding protein that plays a role in the regulation of numerous cellular processes, such as cell differentiation, cell division, cell proliferation, cell migration, thrombosis, angiogenesis, cardiac hypertrophy and apoptosis. Involved in bone marrow megakaryocyte differentiation by negatively regulating thrombopoietin-mediated signaling pathway. Participates in the endomitotic cell cycle of megakaryocyte, a form of mitosis in which both karyokinesis and cytokinesis are interrupted. Plays a role in integrin signaling by negatively regulating alpha-IIb/beta3 activation in thrombin-stimulated megakaryocytes preventing platelet aggregation. Up-regulates PTK2/FAK1 activity, and is also needed for the recruitment of PTK2/FAK1 to focal adhesions; it thus appears to play an important role in focal adhesion formation. Positively regulates cell migration on fibronectin in a CDC42-dependent manner, the effect being negatively regulated by PAK1. Functions as a negative regulator of stress activated MAP kinase (MAPK) signaling pathways. Down-regulates inositol 1,4,5-trisphosphate receptor-dependent calcium signaling. Involved in sphingosine kinase SPHK1 translocation to the plasma membrane in a N-myristoylation-dependent manner preventing TNF-alpha-induced apoptosis. Regulates serine/threonine-protein kinase PLK3 activity for proper completion of cell division progression. Plays a role in microtubule (MT) dynamics during neuronal development; disrupts the MT depolymerization activity of STMN2 attenuating NGF-induced neurite outgrowth and the MT reorganization at the edge of lamellipodia. Promotes cardiomyocyte hypertrophy via activation of the calcineurin/NFAT signaling pathway. Stimulates calcineurin PPP3R1 activity by mediating its anchoring to the sarcolemma. In ischemia-induced (pathological or adaptive) angiogenesis, stimulates endothelial cell proliferation, migration and microvessel formation by activating the PAK1 and ERK1/ERK2 signaling pathway. Promotes also cancer cell survival and proliferation. May regulate cell cycle and differentiation of spermatogenic germ cells, and/or differentiation of supporting Sertoli cells.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) uses an alternate in-frame splice junction at the 5' end of an exon compared to variant 1. The resulting isoform (2) has the same N- and C-termini but is shorter compared to isoform 1.