

Product datasheet for MC226175

Vegfa (NM 001287057) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Vegfa (NM_001287057) Mouse Untagged Clone

Tag: Tag Free
Symbol: Vegfa

Synonyms: V; Veg; Vegf; VEGF12; VEGF16; VEGF18; Vpf

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226175 representing NM_001287057

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001287057

Insert Size: 573 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001287057.1</u>, <u>NP 001273986.1</u>

 RefSeq Size:
 3475 bp

 RefSeq ORF:
 573 bp

 Locus ID:
 22339

 UniProt ID:
 Q00731

 Cytogenetics:
 17 22.79 cM

Gene Summary: This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding

protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site.[provided by RefSeq, Nov 2015] Transcript Variant: This variant (2) lacks an internal, in-frame coding exon compared to variant 1. It can initiate translation from a non-AUG (CUG) codon, and also from an in-frame, downstream AUG start codon. The isoform (Vegf-A) represented in this RefSeq is derived from the AUG start codon. It has a shorter N-terminus and lacks an internal protein segment compared to isoform 1. This variant also encodes a C-terminally extended isoform (Vegf-Ax), which results from the use of an alternative in-frame, downstream translation termination codon (UAA). Isoform Vegf-Ax is represented in NM_001317041.1.