

Product datasheet for MC226205

Csrp3 (NM_001198841) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Csrp3 (NM_001198841) Mouse Untagged Clone

Tag: Tag Free
Symbol: Csrp3

Synonyms: CRP3; MLP; MMLP

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226205 representing NM_001198841

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCCAAACTGGGGTGGAGGTGCAAAATGTGGAGCCTGTGAAAAGACGGTCTACCATGCAGAAGAAATCC
AGTGCAATGGGAGGAGGATTTCCACAAGACCTGTTTCCACTGCATGGCCTGCAGGAAAGCTCTGGACAGCAC
CACAGTGGCAGCTCATGAGTCAGAGATCTACTGTAAGGTGTGCTATGGGCGCAGGTATGGCCCCAAGGGG
ATCGGGTTCGGACAAGGCGCTGGCTGCCTCAGCACAGACACTGGCGAGCATCTTGGCCTGCAGTTCCAAC
AATCCCCAAAGCCAGCTCGAGCAGCCACCACAAGCAACCCTTCCAAATTCTCTGCAAAGTTTGGAGAATC
AGAGAAGTGCCCACGATGTGGAAAGTCGGTATACGCTGCTGAGAAGGTCATGGGAGGTGGCAAGCCCTGG
CACAAGACCTGCTTCCGCTGTGCCATCTGTGGGAAGGCCTGGAGTCTACAAATGTCACTGACAAGGATG
GGGAGCTCTACTGCAAAGTTTGCTATGCCAAAAATTTTGGCCCCACAGGCATTGGGTTTGGAGGGCTTAC
ACAGCAAGTGGAAAAGAAGAAGAAGTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001198841

Insert Size: 585 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Csrp3 (NM_001198841) Mouse Untagged Clone - MC226205

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001198841.1</u>, <u>NP 001185770.1</u>

RefSeq Size: 1035 bp
RefSeq ORF: 585 bp
Locus ID: 13009
UniProt ID: P50462
Cytogenetics: 7 B4

Gene Summary: Positive regulator of myogenesis. Acts as cofactor for myogenic bHLH transcription factors

such as MYOD1, and probably MYOG and MYF6. Enhances the DNA-binding activity of the MYOD1:TCF3 isoform E47 complex and may promote formation of a functional MYOD1:TCF3 isoform E47:MEF2A complex involved in myogenesis (By similarity). Plays a crucial and specific

role in the organization of cytosolic structures in cardiomyocytes. Could play a role in mechanical stretch sensing. May be a scaffold protein that promotes the assembly of interacting proteins at Z-line structures. It is essential for calcineurin anchorage to the Z line.

Required for stress-induced calcineurin-NFAT activation (PubMed:9039266,

PubMed:15665106). The role in regulation of cytoskeleton dynamics by association with CFL2 is reported conflictingly. Proposed to contribute to the maintenance of muscle cell integerity through an actin-based mechanism. Can directly bind to actin filaments, cross-link actin filaments into bundles without polarity selectivity and protect them from dilution- and cofilinmediated depolymerization; the function seems to involve its self-association (By similarity). In vitro can inhibit PKC/PRKCA activity. Proposed to be involved in cardiac stress signaling by down-regulating excessive PKC/PRKCA signaling (PubMed:27353086).[UniProtKB/Swiss-Prot

Function]

Transcript Variant: This variant (2) differs in the 5' UTR, compared to variant 1. Variants 1 and

2 encode the same protein.