

## **Product datasheet for MC226216**

## Inca1 (NM 001252485) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Inca1 (NM\_001252485) Mouse Untagged Clone

Tag: Tag Free
Symbol: Inca1

Synonyms: Al842396

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226216 representing NM\_001252485

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001252485

**Insert Size:** 591 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## Inca1 (NM\_001252485) Mouse Untagged Clone - MC226216

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001252485.1</u>, <u>NP 001239414.1</u>

RefSeq Size: 1309 bp
RefSeq ORF: 591 bp
Locus ID: 103844
UniProt ID: Q6PKN7
Cytogenetics: 11 B3

**Gene Summary:** Binds to CDK2-bound cyclins and inhibits the kinase activity of CDK2; binding to cyclins is

critical for its function as CDK inhibitor. Inhibits cell growth and proliferation and may play a role in cell cycle control (PubMed:21540187). Required for ING5-mediated regulation of Sphase progression, enhancement of Fas-induced apoptosis and inhibition of cell growth

(PubMed:21750715).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) differs in the 5' sequence, which results in a downstream AUG start codon, compared to variant 1. The resulting isoform (2) is shorter at the N-terminus, compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly.

The genomic coordinates used for the transcript record were based on alignments.