

## **Product datasheet for MC226269**

## Chtop (NM 001293779) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Chtop (NM\_001293779) Mouse Untagged Clone

Tag: Tag Free Symbol: Chtop

Synonyms: 2500003M10Rik; Fop; Srag

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC226269 representing NM\_001293779

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCTGCACAGTCAGCGCCGAAAGTTGTGCTAAAAAAGCACCACCAAGATGTCTCTAAATGAGCGCTTTA
CTAATATGCTGAAGAACAAACAGCCGATGCCAGTGAATATTCGGGCTTCGATGCAGCAGCAGCAGCAGCT
AGCCAGTGCCAGAAACAGAAGACTGGCCCAGCAGATGGAGAATAGACCCTCTGTCCAGGCAGCATTAAAA
CTTAAGCAGAGCTTAAAGCAGCGCCTGGGTAAGAGTAATATCCAGGCACGGTTAGGCCGACCCATAGGTG
CCCTGGCCAGGGGAGCAATTGGAGGAAGAGGCCTACCCATAATCCAGAGAGGCTTGCCCCGAGGAGGACT
ACGTGGGGGACGTGCTACCAGAACCCTGCTTAGGGGTGGGATGTCGCTCCGAGGTCGGGTATGATAGGT
CGGGGAAGAGGGGGCTTTGGAGGCAGAGCCCGAGGTCGGCCGAGGAGGAGGTGCCCTCACTCGCCCTG
TATTGACCAAGGAGCAGCTGGACAACCAATTGGATGCATACATGTCGAAAACTAAAAGGACACCTGGATGC

TGAATTGGATGCCTACATGGCACAGACAGATCCTGAAACCAATGAT<mark>TGA</mark>

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_001293779

**Insert Size:** 609 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

> point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: NM 001293779.1, NP 001280708.1

RefSeq Size: 2297 bp RefSeq ORF: 609 bp Locus ID: 66511

**UniProt ID:** Q9CY57 3 F1

Cytogenetics:

**Gene Summary:** Plays an important role in the ligand-dependent activation of estrogen receptor target genes

> (By similarity). May play a role in the silencing of fetal globin genes (PubMed:20688955). Recruits the 5FMC complex to ZNF148, leading to desumoylation of ZNF148 and subsequent transactivation of ZNF148 target genes (PubMed:22872859). Required for the tumorigenicity of glioblastoma cells. Binds to 5-hydroxymethylcytosine (5hmC) and associates with the methylosome complex containing PRMT1, PRMT5, MEP50 and ERH. The CHTOP-methylosome

complex associated with 5hmC methylates H4R3 and transactivates genes involved in

glioblastomagenesis (PubMed:25284789).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (5) uses an alternate in-frame splice site in the 5' coding region and lacks an in-frame exon in the 3' coding region compared to variant 1. The encoded isoform (5) is shorter than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on

transcript alignments.