

## **Product datasheet for MC226277**

## Cited1 (NM\_001276473) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Cited1 (NM\_001276473) Mouse Untagged Clone

Tag: Tag Free
Symbol: Cited1

**Synonyms:** Al316840; AU019144; Msg1

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC226277 representing NM\_001276473

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGCCAACTATGTCGAGGCCTGCACTTGATGTCAAGGGTGGCACCACCTCTGGGAAGGAGGATGCCAACC
AGGAGATGAACTCTCTGGCCTACTCCAACCTTGGAGTGAAGGATCGCAAGGCAGTGACTGTCCTGCACTA
CCCCGGGGTCACCGCAAATGGAGCCAAAGCCAACGGAGTTCCCACTAGCTCCTCTGGATCGACATCTCCA
ATAGGCTCTCCTACTGCCACCCCTTCTTCCAAACCCCCATCCTTCAACCTGCATCCTACCCCTCACCTGA
TGGCCAGCATGCAGCTTCAGAAGCTTAATAGCCAGTACCAAGGGGCTGCGGCTACTGCTGCTGCTCCT
CACTGGTGCAGGCCTACCAGGGGAGGAAGAGCCCATGCAAAACTGGGTCACCGCCCCTCTGGTAGTGGGA
GGGTCTCCGGGATCTGTCTCCTCCTGCTGGTGGCCCAGAGCCCTGCTCCATTGATTCTGACCCGGTGG
ATGAGGGGTGCTGATGTCTCTCTGTGGTGTTGAATTGGGCCTAGACCGAGCCCAATGAGCTTCCCGAGCTGTG

 ${\tt GCTGGGGCAGAATGAGTTTGATTTCACTGCAGATTTTCCCTCTGGCTGC} {\tt TGA}$ 

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_001276473

**Insert Size:** 612 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

## Cited1 (NM\_001276473) Mouse Untagged Clone - MC226277

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001276473.1</u>, <u>NP 001263402.1</u>

X 45.25 cM

 RefSeq Size:
 773 bp

 RefSeq ORF:
 612 bp

 Locus ID:
 12705

 UniProt ID:
 P97769

Cytogenetics:

**Gene Summary:** Transcriptional coactivator of the p300/CBP-mediated transcription complex. Enhances

SMAD-mediated transcription by strengthening the functional link between the DNA-binding SMAD transcription factors and the p300/CBP transcription coactivator complex. Stimulates estrogen-dependent transactivation activity mediated by estrogen receptors signaling; stabilizes the interaction of estrogen receptor ESR1 and histone acetyltransferase EP300. Positively regulates TGF-beta signaling through its association with the SMAD/p300/CBP-mediated transcriptional coactivator complex. Induces transcription from estrogen-responsive promoters and protection against cell death. Potentiates EGR2-mediated transcriptional activation activity from the ERBB2 promoter. Acts as an inhibitor of osteoblastic mineralization through a cAMP-dependent parathyroid hormone receptor signaling. May play a role in pigmentation of melanocytes. Associates with chromatin to the

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) differs in the 5' UTR, compared to variant 1. Variants 1, 2

estrogen-responsive TGF-alpha promoter region in a estrogen-dependent manner.

and 3 encode the same isoform (a).