

## Product datasheet for MC226309

### Med27 (NM\_029615) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Med27 (NM\_029615) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Med27  
**Synonyms:** 1500015J03Rik; 2310042P07Rik; AA682045; Crsp8; D2Ertd434e  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >MC226309 representing NM\_029615  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGC**C

ATGGCGGACGTGCTGAGCGTTGGTGTGAACCTGGAGGCCTTTTCGCAGGCTATTAGCGCCATCCAGGCGC  
 TACGCTCTAGCGTGAGTAGGGTCTTCGACTGCCTTAAGGATGGCATGCGCAACAAAGAGACGCTGGAGGG  
 CCGCGAGAAGGCCTTCATCGCGAACTCCAGGACAACCTGCATTCCGTCACCGGGACCTCAATGAACTG  
 GAACGTTTGAGCAACCTGGTAGGCAAGCCGTCTGAGAACCACCCTTTCATAACAGTGGGCTGCTAAGCC  
 TGGATCCTGTGCAGGACAAAACCCCACTCTACAGCCAGCTTCTGCAAGCATACAAATGGTCCAACAAGCT  
 GCAGTACCATGCAGGTCTAGCGTCTGGCCTTTTGAACCAACAGTCCTTGAAGCGCTCTGCAAATCAGATG  
 GGTGTGTCTGCCAAGCGGAGACCGAAGGCCAGCCCACTCTGTTTCTACCACCTCAGTATGTCGATG  
 ACGTGATCAGCCGATTGACAGAATGTTTCCAGAGATGTCCATTCACTTATCGAGACCAATGGGACTTC  
 AGCAATGCTTCTGATTCTTATAACAGCTCTGCTCAGAGCATTACTGGCCCTTGGTTTGGTCT**GA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-MluI  
**ACCN:** NM\_029615  
**Insert Size:** 627 bp


[View online »](#)

<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>OTI Annotation:</b>	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_029615.1</a> , <a href="#">NP_083891.1</a>
<b>RefSeq Size:</b>	1709 bp
<b>RefSeq ORF:</b>	627 bp
<b>Locus ID:</b>	68975
<b>UniProt ID:</b>	<a href="#">Q9DB40</a>
<b>Cytogenetics:</b>	2 19.84 cM
<b>Gene Summary:</b>	<p>Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors (By similarity). [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) lacks four 3' exons but instead contains an alternate 3' terminal exon, and it thus differs in the 3' coding region and 3' UTR, compared to variant 1. The encoded isoform (b) has a distinct C-terminus and is shorter than isoform a.</p>