

Product datasheet for **MC226395**

Lair1 (NM_001302675) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Lair1 (NM_001302675) Mouse Untagged Clone
Tag: Tag Free
Symbol: Lair1
Synonyms: 5133400O11Rik; BB115266; D7Bwg0421e; Lair-1
Mammalian Cell Selection: Neomycin
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Fully Sequenced ORF: >MC226395 representing NM_001302675
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGTCACCTTCATCCAGTTATCCTGCTGGTCTTGTGCTGTGCCTGGGATGGAAAATTAACACACAGGAGG
 GTTCTCTGCCTGATATTACCATCTTCCCTAATTCAAGTCTTATGATCTCCAAGGGACTTTTGTAACTGT
 TGTGTGCTCATACTCTGATAAACACGACTTGTATAACATGGTCCGCTGGAGAAGGACGGCAGCACCTTT
 ATGGAAAAGAGCACTGAGCCTTATAAACAGAGGATGAATTTGAGATTGGCCAGTGAATGAAACCATTA
 CTGGACATTATAGCTGTATCTATTCTGAAGGGGATTACCTGGTCCGAACGTAGTAAGACGCTGGAGCTAAA
 GGTGATCAAAGAAAATGTCATCCAGACTCTGCCCCAGGTCCAACCTCAGGACTCCCAAACAACAAAAGA
 CAGCAGCAGAGGCCAGAAGAGAGGCTAAATCTAGCTACTAATGGCCTGGAGATGACTCCAGACATAGTTG
 CAGATGACAGGCTTCTGAGGACAGATGGACAGAACTGGACCCAGTTGCAGGAGACCTTCAAGAGGT
 GACGTATATCCAGCTGGACCATCACTCCCTCACACAGAGGGCAGTCGGAGCTGTGACCTCACAGAGCACA
 GATATGGCTGAGTCCAGCACATATGCAGCCATCATCAGACACT**GTA**

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001302675
Insert Size: 675 bp



[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_001302675.1</u> , <u>NP_001289604.1</u>
RefSeq Size:	3395 bp
RefSeq ORF:	675 bp
Locus ID:	52855
UniProt ID:	<u>Q8BG84</u>
Cytogenetics:	7 2.31 cM
Gene Summary:	<p>Functions as an inhibitory receptor that plays a constitutive negative regulatory role on cytolytic function of natural killer (NK) cells, B-cells and T-cells. Activation by Tyr phosphorylation results in recruitment and activation of the phosphatases PTPN6 and PTPN11. It also reduces the increase of intracellular calcium evoked by B-cell receptor ligation. May also play its inhibitory role independently of SH2-containing phosphatases. Modulates cytokine production in CD4+ T-cells, down-regulating IL2 and IFNG production while inducing secretion of transforming growth factor beta. Down-regulates also IgG and IgE production in B-cells as well as IL8, IL10 and TNF secretion. Inhibits proliferation and induces apoptosis in myeloid leukemia cell lines as well as prevents nuclear translocation of NF-kappa-B p65 subunit/RELA and phosphorylation of I-kappa-B alpha/CHUK in these cells. Inhibits the differentiation of peripheral blood precursors towards dendritic cells (By similarity).[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (e) lacks an in-frame exon in the coding region compared to variant a. The encoded isoform (e) is shorter than isoform a.</p>