

Product datasheet for MC226497

Derl2 (NM 001291148) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Derl2 (NM_001291148) Mouse Untagged Clone

Tag: Tag Free Symbol: Derl2

Synonyms: CGI-101; Derlin-2; F-lana; Flana

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226497 representing NM_001291148

Red=Cloning site Blue=ORF Orange=Stop codon

 ${\tt TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCCC}$

GCCGCGATCGCC

AGCGCCTTGGTGGGTAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM 001291148

Insert Size: 717 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Derl2 (NM_001291148) Mouse Untagged Clone - MC226497

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001291148.1</u>, <u>NP 001278077.1</u>

 RefSeq Size:
 3665 bp

 RefSeq ORF:
 717 bp

 Locus ID:
 116891

UniProt ID: Q8BNI4

Cytogenetics: 11 43.21 cM

Gene Summary: Functional component of endoplasmic reticulum-associated degradation (ERAD) for

misfolded lumenal glycoproteins, but not that of misfolded nonglycoproteins. May act by forming a channel that allows the retrotranslocation of misfolded glycoproteins into the cytosol where they are ubiquitinated and degraded by the proteasome. May mediate the interaction between VCP and misfolded glycoproteins. May also be involved in endoplasmic reticulum stress-induced pre-emptive quality control, a mechanism that selectively attenuates the translocation of newly synthesized proteins into the endoplasmic reticulum and reroutes

them to the cytosol for proteasomal degradation.[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (4) uses an alternate in-frame splice site in the 5' coding region, compared to variant 1. The encoded isoform (c) is shorter, compared to isoform a.