

Product datasheet for MC226531

Dcun1d1 (NM_001205362) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Dcun1d1 (NM 001205362) Mouse Untagged Clone

Tag: Tag Free Symbol: Dcun1d1

Synonyms: pTes3; Rp42; SCCRO; Tes3

Mammalian Cell N

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC226531 representing NM_001205362

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001205362

Insert Size: 735 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Dcun1d1 (NM_001205362) Mouse Untagged Clone - MC226531

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 001205362.1</u>, <u>NP 001192291.1</u>

 RefSeq Size:
 4434 bp

 RefSeq ORF:
 735 bp

 Locus ID:
 114893

Cytogenetics: 3 B

Gene Summary: Part of an E3 ubiquitin ligase complex for neddylation. Promotes neddylation of cullin

components of E3 cullin-RING ubiquitin ligase complexes. Acts by binding to cullin-RBX1 complexes in the cytoplasm and promoting their nuclear translocation, enhancing recruitment of E2-NEDD8 (UBE2M-NEDD8) thioester to the complex, and optimizing the orientation of proteins in the complex to allow efficient transfer of NEDD8 from the E2 to the cullin substrates. Involved in the release of inhibitory effets of CAND1 on cullin-RING ligase E3 complex assembly and activity (By similarity). Acts also as an oncogene facilitating malignant transformation and carcinogenic in vivo (PubMed:20563250).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) differs in the 5' UTR, lacks a portion of the 5' coding region and uses a downstream, in-frame start codon, compared to variant 1. The encoded isoform (2) has a shorter N-terminus, compared to isoform 1. Variants 2 and 3 encode the same isoform (2). Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.