

## **Product datasheet for MC226552**

## Gsto2 (NM 030051) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Gsto2 (NM\_030051) Mouse Untagged Clone

Tag: Tag Free
Symbol: Gsto2

**Synonyms:** 1700020F09Rik; 4930425C18Rik; GSTO 2-2

Mammalian Cell

Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_030051

**Insert Size:** 747 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>NM 030051.1</u>, <u>NP 084327.1</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Gsto2 (NM\_030051) Mouse Untagged Clone - MC226552

 RefSeq Size:
 1258 bp

 RefSeq ORF:
 747 bp

 Locus ID:
 68214

 UniProt ID:
 Q8K2Q2

 Cytogenetics:
 19 D1

**Gene Summary:** Exhibits glutathione-dependent thiol transferase activity. Has high dehydroascorbate

reductase activity and may contribute to the recycling of ascorbic acid. Participates in the biotransformation of inorganic arsenic and reduces monomethylarsonic acid (MMA).

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (2) differs in the 5' UTR, compared to variant 1. Variants 1 and

2 encode the same protein.