

## **Product datasheet for MC226583**

## Mdfi (NM\_001276391) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Mdfi (NM\_001276391) Mouse Untagged Clone

Tag: Tag Free
Symbol: Mdfi

Synonyms: I-mf; I-mfa
Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC226583 representing NM\_001276391

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul

**ACCN:** NM\_001276391

**Insert Size:** 756 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com **OTI Disclaimer:** 

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**OTI Annotation:** Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube Components:

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

NM 001276391.1, NP 001263320.1 RefSeq:

RefSeq Size: 1483 bp RefSeq ORF: 756 bp Locus ID: 17240

Cytogenetics: 17 23.99 cM

**Gene Summary:** Inhibits the transactivation activity of the Myod family of myogenic factors and represses

> myogenesis. Acts by associating with Myod family members and retaining them in the cytoplasm by masking their nuclear localization signals. Can also interfere with the DNAbinding activity of Myod family members. Plays an important role in trophoblast and chondrogenic differentiation. Regulates the transcriptional activity of TCF7L1/TCF3 by interacting directly with TCF7L1/TCF3 and preventing it from binding DNA. Binds to the axin complex, resulting in an increase in the level of free beta-catenin. Affects axin regulation of

the WNT and JNK signaling pathways.[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4) uses an alternate 5' exon structure, differs in the 5' UTR, contains an alternate exon in the 5' coding region and uses a downstream start codon compared to variant 2. It encodes isoform 2 which is longer and has a distinct N-terminus,

compared to isoform 1.