

Product datasheet for MC226629

Cfd (NM_001291915) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cfd (NM_001291915) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Cfd
Synonyms:	A; Adn; DF
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001291915
Insert Size:	777 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
OTI Annotation:	Clone contains native stop codon, and expresses the complete ORF without any c-terminal tag.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.


[View online »](#)

RefSeq: [NM_001291915.2](#), [NP_001278844.1](#)

RefSeq Size: 939 bp

RefSeq ORF: 777 bp

Locus ID: 11537

UniProt ID: [P03953](#)

Cytogenetics: 10 39.72 cM

Gene Summary: This gene encodes a serine protease that plays an important role in the alternative pathway of complement activation for pathogen recognition and elimination. The encoded preproprotein undergoes proteolytic processing to generate a mature, functional enzyme that in turn cleaves factor B in the complement pathway. This gene is expressed in adipocytes and the mature enzyme is secreted into the bloodstream. Mice lacking the encoded product cannot initiate alternative pathway of complement activation. [provided by RefSeq, Jul 2016]
Transcript Variant: This variant (2) uses an alternate in-frame splice site in the 5' coding region, compared to variant 1. The encoded protein (isoform 2) is shorter, compared to isoform 1.