

Product datasheet for MC226638

Nudt7 (NM_001290181) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nudt7 (NM_001290181) Mouse Untagged Clone

Tag: Tag Free Symbol: Nudt7

Synonyms: 1300007B24Rik; 2210404C19Rik

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC226638 representing NM_001290181

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001290181

Insert Size: 783 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com **OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

OTI Annotation: Clone contains native stop codon, and expresses the complete ORF without any c-terminal

tag.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001290181.1, NP 001277110.1</u>

 RefSeq Size:
 2839 bp

 RefSeq ORF:
 783 bp

 Locus ID:
 67528

 UniProt ID:
 099P30

Cytogenetics: 8 E1

Gene Summary: Coenzyme A diphosphatase which mediates the cleavage of CoA, CoA esters and oxidized

CoA with similar efficiencies, yielding 3',5'-ADP and the corresponding 4'-phosphopantetheine derivative as products. CoA into 3',5'-ADP and 4'-phosphopantetheine. Has no activity toward

NDP-sugars, CDP-alcohols, (deoxy)nucleoside 5'-triphosphates, nucleoside 5'-di or

monophosphates, diadenosine polyphosphates, NAD, NADH, NADP, NADPH or thymidine-5'-

monophospho-p-nitrophenyl ester. May be required to eliminate oxidized CoA from

peroxisomes, or regulate CoA and acyl-CoA levels in this organelle in response to metabolic

demand. Does not play a role in U8 snoRNA decapping activity. Binds U8 snoRNA.

[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (4) contains an alternate 5' terminal exon, and it thus differs in the 5' UTR and initiates translation at an alternate start codon, compared to variant 1. The encoded isoform (4) has a distinct N-terminus and is longer than isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.